




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07C 311/39, A61K 31/18, A61P 5/48, 17/06, 19/02, 25/28, 31/18, 35/00, 37/06		A1	(11) International Publication Number: WO 00/42003
			(43) International Publication Date: 20 July 2000 (20.07.00)
(21) International Application Number: PCT/US99/30435 (22) International Filing Date: 21 December 1999 (21.12.99) (30) Priority Data: 60/115,874 13 January 1999 (13.01.99) US 60/122,422 2 March 1999 (02.03.99) US (71) Applicant (for all designated States except US): WARNER-LAMBERT COMPANY [US/US]; 201 Tabor Road, Morris Plains, NJ 07950 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): BARRETT, Stephen, Douglas [US/US]; 14220 Sunbury, Livonia, MI 48154 (US). TECLE, Haile [US/US]; 3048 Tumberry, Ann Arbor, MI 48108 (US). BOOTH, Richard, John [GB/US]; 1433 Natalie Lane, Apt. 104, Ann Arbor, MI 48105 (US). (74) Agents: RYAN, M., Andrea; Warner-Lambert Company, 201 Tabor Road, Morris Plains, NJ 07950 (US) et al.			(81) Designated States: AE, AL, AU, BA, BB, BG, BR, CA, CN, CR, CU, CZ, DM, EE, GD, GE, HR, HU, ID, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MA, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ZA, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>

(54) Title: BENZENESULFONAMIDE DERIVATIVES AND THEIR USE AS MEK INHIBITORS



(57) Abstract

Benzenesulfonamides of formula (I), in which W is OR₁, NR₂OR₁, NR_AR_B, NR₂NR_AR_B, or NR₂(CH₂)₂₋₄NR_AR_B and the other variables as defined in the claims, are inhibitors of MEK and are effective in the treatment of proliferative diseases, cancer, stroke, heart failure, xenograft rejection, arthritis, cystic fibrosis, hepatomegaly, cardiomegaly, Alzheimer's disease, complications of diabetes, septic shock, and viral infection.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

BENZENESULFONAMIDE DERIVATIVES AND THEIR USE AS MEK INHIBITORS

The invention relates to benzenesulfonamides and derivatives thereof.

5

BACKGROUND

MEK enzymes are dual specificity kinases involved in, for example, immunomodulation, inflammation, and proliferative diseases such as cancer and restenosis.


10 Proliferative diseases are caused by a defect in the intracellular signaling system, or the signal transduction mechanism of certain proteins. Defects include a change either in the intrinsic activity or in the cellular concentration of one or more signaling proteins in the signaling cascade. The cell may produce a growth factor that binds to its own receptors, resulting in an autocrine loop, which
15 continually stimulates proliferation. Mutations or overexpression of intracellular signaling proteins can lead to spurious mitogenic signals within the cell. Some of the most common mutations occur in genes encoding the protein known as Ras, a G-protein that is activated when bound to GTP, and inactivated when bound to GDP. The above-mentioned growth factor receptors, and many other mitogenic
20 receptors, when activated, lead to Ras being converted from the GDP-bound state to the GTP-bound state. This signal is an absolute prerequisite for proliferation in most cell types. Defects in this signaling system, especially in the deactivation of the Ras-GTP complex, are common in cancers, and lead to the signaling cascade below Ras being chronically activated.

25 Activated Ras leads in turn to the activation of a cascade of serine/threonine kinases. One of the groups of kinases known to require an active Ras-GTP for its own activation is the Raf family. These in turn activate MEK (e.g., MEK₁ and MEK₂) which then activates MAP kinase, ERK (ERK₁ and ERK₂). Activation of MAP kinase by mitogens appears to be essential for
30 proliferation; constitutive activation of this kinase is sufficient to induce cellular transformation. Blockade of downstream Ras signaling, for example by use of a dominant negative Raf-1 protein, can completely inhibit mitogenesis, whether

induced from cell surface receptors or from oncogenic Ras mutants. Although Ras is not itself a protein kinase, it participates in the activation of Raf and other kinases, most likely through a phosphorylation mechanism. Once activated, Raf and other kinases phosphorylate MEK on two closely adjacent serine residues, S²¹⁸ and S²²² in the case of MEK-1, which are the prerequisite for activation of MEK as a kinase. MEK in turn phosphorylates MAP kinase on both a tyrosine, Y¹⁸⁵, and a threonine residue, T¹⁸³, separated by a single amino acid. This double phosphorylation activates MAP kinase at least 100-fold. Activated MAP kinase can then catalyze the phosphorylation of a large number of proteins, including several transcription factors and other kinases. Many of these MAP kinase phosphorylations are mitogenically activating for the target protein, such as a kinase, a transcription factor, or another cellular protein. In addition to Raf-1 and MEKK, other kinases activate MEK, and MEK itself appears to be a signal integrating kinase. Current understanding is that MEK is highly specific for the phosphorylation of MAP kinase. In fact, no substrate for MEK other than the MAP kinase, ERK, has been demonstrated to date and MEK does not phosphorylate peptides based on the MAP kinase phosphorylation sequence, or even phosphorylate denatured MAP kinase. MEK also appears to associate strongly with MAP kinase prior to phosphorylating it, suggesting that phosphorylation of MAP kinase by MEK may require a prior strong interaction between the two proteins. Both this requirement and the unusual specificity of MEK are suggestive that it may have enough difference in its mechanism of action to other protein kinases that selective inhibitors of MEK, possibly operating through allosteric mechanisms rather than through the usual blockade of the ATP binding site, may be found.

SUMMARY

The invention features a compound having the formula (I) below:



(I)

- W is OR₁, NR₂OR₁, NR_AR_B, NR₂NR_AR_B, or NR₂(CH₂)₂₋₄NR_AR_B. R₁ is H, C₁₋₈ alkyl, C₃₋₈ alkenyl, C₃₋₈ alkynyl, C₃₋₈ cycloalkyl, phenyl, (phenyl)C₁₋₄ alkyl, (phenyl)C₃₋₄ alkenyl, (phenyl)C₃₋₄ alkynyl, (C₃₋₈ cycloalkyl)-C₁₋₄ alkyl, (C₃₋₈ cycloalkyl)C₃₋₄ alkenyl, (C₃₋₈ cycloalkyl)C₃₋₄ alkynyl, C₃₋₈ heterocyclic radical, (C₃₋₈ heterocyclic radical)C₁₋₄ alkyl, (C₃₋₈ heterocyclic radical)C₃₋₄ alkenyl, (C₃₋₈ heterocyclic radical)C₃₋₄ alkynyl, or (CH₂)₂₋₄NR_AR_B. R₂ is H, phenyl, C₁₋₄ alkyl, C₃₋₄ alkenyl, C₃₋₈ alkynyl, C₃₋₈ cycloalkyl, or (C₃₋₈ cycloalkyl)C₁₋₄ alkyl. R_A is H, C₁₋₆ alkyl, C₃₋₈ alkenyl, C₃₋₈ alkynyl, C₃₋₈ cycloalkyl, phenyl, (C₃₋₈ cycloalkyl)C₁₋₄ alkyl, (C₃₋₈ cycloalkyl)C₃₋₄ alkenyl, (C₃₋₈ cycloalkyl)C₃₋₄ alkynyl, C₃₋₈ heterocyclic radical, (C₃₋₈ heterocyclic radical)C₁₋₄ alkyl, (aminosulfonyl)phenyl, [(aminosulfonyl)phenyl]C₁₋₄ alkyl, (aminosulfonyl)C₁₋₆ alkyl, (aminosulfonyl)C₃₋₆ cycloalkyl, or [(aminosulfonyl)C₃₋₆ cycloalkyl]C₁₋₄ alkyl. R_B is H, C₁₋₈ alkyl, C₃₋₈ alkenyl, C₃₋₈ alkynyl, C₃₋₈ cycloalkyl, or C₆₋₈ aryl. R₃ is H, F, Cl, Br, or NO₂. R₄ is H or F. R₅ is H, methyl or Cl. R₆ is H, C₁₋₄ alkyl, hydroxyethyl, hydroxypropyl, (CH₂)₂₋₄(NR_CR_D), phenyl, 2-pyridyl, 3-pyridyl, 4-pyridyl or CH₂Ar, where Ar is phenyl, 2-pyridyl, 3-pyridyl, or 4-pyridyl. R₇ is H, C₁₋₄ alkyl, hydroxyethyl, hydroxypropyl, (CH₂)₂₋₄(NR_CR_D), phenyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, or CH₂Ar', where Ar' is phenyl, 2-pyridyl, 3-pyridyl, or 4-pyridyl. Each of R_C and R_D is independently selected from H, C₁₋₆ alkyl, C₃₋₄ alkenyl, C₃₋₄ alkynyl, C₃₋₆ cycloalkyl, C₃₋₆ heterocyclic radical, and phenyl. NR_CR_D can also be selected from morpholinyl, piperazinyl, pyrrolidinyl, or piperadinyl. Each

hydrocarbon radical or heterocyclic radical above is optionally substituted with between 1 and 3 substituents independently selected from halo, C₁₋₄ alkyl, C₃₋₆ cycloalkyl, C₂₋₄ alkenyl, C₂₋₄ alkynyl, phenyl, hydroxy, amino, (amino)sulfonyl, and NO₂, wherein each substituent alkyl, cycloalkyl, alkenyl, alkynyl or phenyl is
 5 in turn optionally substituted with between 1 and 3 substituents independently selected from halo, C₁₋₂ alkyl, hydroxy, amino, and NO₂. The invention also features pharmaceutically acceptable salts and C₁₋₇ esters thereof.

Preferred compounds include PD 297764, 3,4-Difluoro-2-(4-iodo-phenylamino)-N-methoxy-5-(4-pyridin-2-yl-piperazine-1-sulfonyl)-benzamide;
 10 PD 297765, N-Allyloxy-3,4-difluoro-2-(4-iodo-phenylamino)-5-(4-methyl-piperazine-1-sulfonyl)-benzamide; PD297766, N-Allyloxy-5-[(2-diethylamino-ethyl)-methyl-sulfamoyl]-3,4-difluoro-2-(4-iodo-phenylamino)-benzamide;
 PD297767, N-Allyloxy-5-[(3-dimethylamino-propyl)-methyl-sulfamoyl]-3,4-difluoro-2-(4-iodo-phenylamino)-benzamide; PD297768, N-Cyclopropylmethoxy-3,4-
 15 difluoro-2-(4-iodo-phenylamino)-5-(4-methyl-piperazine-1-sulfonyl)-benzamide; PD297769, N-Cyclopropylmethoxy-5-[(2-diethylamino-ethyl)-methyl-sulfamoyl]-3,4-difluoro-2-(4-iodo-phenylamino)-benzamide; PD297770, N-Cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-phenylamino)-5-[methyl-(2-pyridin-2-yl-ethyl)-sulfamoyl]-benzamide; PD297771, N-Cyclopropylmethoxy-3,4-difluoro-2-
 20 (4-iodo-phenylamino)-5-(4-pyridin-2-yl-piperazine-1-sulfonyl)-benzamide; PD297772, 5-[Benzyl-(2-dimethylamino-ethyl)-sulfamoyl]-N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-phenylamino)-benzamide; PD297773, 3,4-Difluoro-2-(4-iodo-2-methyl-phenylamino)-N-methoxy-5-(4-pyridin-2-yl-piperazine-1-sulfonyl)-benzamide; and PD297774, 1-[5-Allyloxycarbamoyl-2,3-difluoro-4-(4-iodo-2-
 25 methyl-phenylamino)-benzenesulfonyl]-piperidine-3-carboxylic acid amide.

The invention also relates to a pharmaceutical composition including
 (a) a compound of formula (I) and (b) a pharmaceutically-acceptable carrier.

The invention further relates to a method for treating proliferative diseases, such as cancer, restenosis, psoriasis, autoimmune disease, and atherosclerosis.

30 Other aspects of the invention include methods for treating MEK-related (including ras-related) cancers, whether solid or hematopoietic. Examples of cancers include colorectal, cervical, breast, ovarian, brain, acute leukemia,

gastric, non-small cell lung, pancreatic and renal cancer. Further aspects of the invention include methods for treating or reducing the symptoms of xenograft (cell(s), skin, limb, organ or bone marrow transplant) rejection, osteoarthritis, rheumatoid arthritis, cystic fibrosis, complications of diabetes (including diabetic retinopathy and diabetic nephropathy), hepatomegaly, cardiomegaly, stroke (such as acute focal ischemic stroke and global cerebral ischemia), heart failure, septic shock, asthma, and Alzheimer's disease. Compounds of the invention are also useful as antiviral agents for treating viral infections such as HIV, hepatitis B virus (HBV), human papilloma virus (HPV), cytomegalovirus (CMV), and Epstein-Barr virus (EBV). These methods include the step of administering to a patient in need of such treatment, or suffering from such a disease or condition, a pharmaceutically-effective amount of a disclosed compound or pharmaceutical composition thereof.

The invention also features methods of combination therapy, such as a method for treating cancer, wherein the method further includes providing radiation therapy or chemotherapy, for example, with mitotic inhibitors such as a taxane or a vinca alkaloid. Examples of mitotic inhibitors include paclitaxel, docetaxel, vincristine, vinblastine, vinorelbine, and vinflunine. Other therapeutic combinations include a MEK inhibitor of the invention and an anticancer agent such as cisplatin, 5-fluorouracil or 5-fluoro-2-4(1H,3H)-pyrimidinedione (5FU), flutamide, and gemcitabine. The chemotherapy or radiation therapy may be administered before, concurrently, or after the administration of a disclosed compound according to the needs of the patient.

The invention also features synthetic methods and synthetic intermediates disclosed herein.

Other aspects of the invention are provided in the description, examples, and claims below.

DETAILED DESCRIPTION

The invention features benzenesulfonamide compounds, pharmaceutical compositions thereof, and methods of using such compounds and compositions.

According to one aspect of the invention, the compounds are MEK inhibitors. MEK inhibition assays include the cascade assay for inhibitors of MAP kinase pathway described at column 6, line 36 to column 7, line 4 of U.S. Patent Number 5,525,625 and the *in vitro* MEK assay at column 7, lines 4-27 of the same patent, the entire disclosure of which is incorporated by reference (see also Examples 5-10 below).

A. Terms

Certain terms are defined below and by their usage throughout this disclosure.

Alkyl groups include aliphatic (i.e., hydrocarbyl or hydrocarbon radical structures containing hydrogen and carbon atoms) with a free valence. Alkyl groups are understood to include straight chain and branched structures. Examples include methyl, ethyl, propyl, isopropyl, butyl, n-butyl, isobutyl, t-butyl, pentyl, isopentyl, 2,3-dimethylpropyl, hexyl, 2,3-dimethylhexyl, 1,1-dimethylpentyl, heptyl, and octyl. Cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl.

Alkyl groups can be substituted with 1, 2, 3 or more substituents which are independently selected from halo (fluoro, chloro, bromo, or iodo), hydroxy, amino, alkoxy, alkylamino, dialkylamino, cycloalkyl, aryl, aryloxy, arylalkyloxy, heterocyclic radical, and (heterocyclic radical)oxy. Specific examples include fluoromethyl, hydroxyethyl, 2,3-dihydroxyethyl, (2- or 3-furanyl)methyl, cyclopropylmethyl, benzyloxyethyl, (3-pyridinyl)methyl, (2- or 3-furanyl)methyl, (2-thienyl)ethyl, hydroxypropyl, aminocyclohexyl, 2-dimethylaminobutyl, methoxymethyl, *N*-pyridinylethyl, diethylaminoethyl, and cyclobutylmethyl.

Alkenyl groups are analogous to alkyl groups, but have at least one double bond (two adjacent sp^2 carbon atoms). Depending on the placement of a double bond and substituents, if any, the geometry of the double bond may be *entgegen* (E), or *zusammen* (Z), *cis*, or *trans*. Similarly, alkynyl groups have at least one triple bond (two adjacent sp carbon atoms). Unsaturated alkenyl or alkynyl groups may have one or more double or triple bonds, respectively, or a mixture thereof; like alkyl groups, unsaturated groups may be straight chain or branched,

and they may be substituted as described both above for alkyl groups and throughout the disclosure by example. Examples of alkenyls, alkynyls, and substituted forms include cis-2-butenyl, trans-2-butenyl, 3-butylnyl, 3-phenyl-2-propynyl, 3-(2'-fluorophenyl)-2-propynyl, 3-methyl(5-phenyl)-4-pentylnyl, 2-hydroxy-2-propynyl, 2-methyl-2-propynyl, 2-propenyl, 4-hydroxy-3-butylnyl, 3-(3-fluorophenyl)-2-propynyl, and 2-methyl-2-propenyl.

In formula (I), alkenyls and alkynyls can be C₂₋₄ or C₂₋₈, and are preferably C₃₋₄ or C₃₋₈.

More general forms of substituted hydrocarbon radicals include hydroxyalkyl, hydroxyalkenyl, hydroxyalkynyl, hydroxycycloalkyl, hydroxyaryl, and corresponding forms for the prefixes amino-, halo- (e.g., fluoro-, chloro-, or bromo-), nitro-, alkyl-, phenyl-, cycloalkyl- and so on, or combinations of substituents. According to formula (I), therefore, substituted alkyls include hydroxyalkyl, aminoalkyl, nitroalkyl, haloalkyl, alkylalkyl (branched alkyls, such as methylpentyl), (cycloalkyl)alkyl, phenylalkyl, alkoxy, alkylaminoalkyl, dialkylaminoalkyl, arylalkyl, aryloxyalkyl, arylalkyloxyalkyl, (heterocyclic radical)alkyl, and (heterocyclic radical)oxyalkyl. R₁ thus includes hydroxyalkyl, hydroxyalkenyl, hydroxyalkynyl, hydroxycycloalkyl, hydroxyaryl, aminoalkyl, aminoalkenyl, aminoalkynyl, aminocycloalkyl, aminoaryl, alkylalkenyl, (alkylaryl)alkyl, (haloaryl)alkyl, (hydroxyaryl)alkynyl, and so forth. Similarly, R_A includes hydroxyalkyl and aminoaryl, and R_B includes hydroxyalkyl, aminoalkyl, and hydroxyalkyl(heterocyclic radical)alkyl.

Heterocyclic radicals, which include but are not limited to heteroaryls, include: furyl, oxazolyl, isoxazolyl, thiophenyl, thiazolyl, pyrrolyl, imidazolyl, 1,3,4-triazolyl, tetrazolyl, pyridinyl, pyrimidinyl, pyridazinyl, indolyl, and their nonaromatic counterparts. Further examples of heterocyclic radicals include piperidyl, quinolyl, isothiazolyl, piperidinyl, morpholynyl, piperazinyl, tetrahydrofuryl, tetrahydropyrrolyl, pyrrolidinyl, octahydroindolyl, octahydrobenzothiofuranyl, and octahydrobenzofuranyl.

Selective MEK 1 or MEK 2 inhibitors are those compounds which inhibit the MEK 1 or MEK 2 enzymes, respectively, without substantially inhibiting other enzymes such as MKK3, PKC, Cdk2A, phosphorylase kinase, EGF, and PDGF

receptor kinases, and C-src. In general, a selective MEK 1 or MEK 2 inhibitor has an IC_{50} for MEK 1 or MEK 2 that is at least one-fiftieth (1/50) that of its IC_{50} for one of the above-named other enzymes. Preferably, a selective inhibitor has an IC_{50} that is at least 1/100, more preferably 1/500, and even more preferably
 5 1/1000, 1/5000, or less than that of its IC_{50} or one or more of the above-named enzymes.

B. Compounds

One aspect of the invention features disclosed compounds shown in
 10 formula (I) in the Summary section.

Examples of compounds of formula (I) have structures wherein:

- (a) the sulfamoyl group is *meta* to W(CO)- and *para* to the bridging NH; (b) the sulfamoyl group is *para* to W(CO)- and *meta* to the bridging NH; (c) R_4 is fluoro;
- (d) R_3 is fluoro; (e) R_3 is H; (f) W is OH; (g) W is NR_2OR_1 ; (h) each of R_3 and R_4 is
 15 fluoro; (i) R_1 has at least one hydroxy substituent; (k) R_1 is H, methyl, ethyl, propyl, isopropyl, isobutyl, benzyl, phenethyl, allyl, C_{2-5} alkenyl, C_{2-5} alkynyl, C_{3-6} cycloalkyl, (C_{3-5} cycloalkyl) C_{1-2} alkyl, (C_{3-5} heterocyclic radical)- C_{1-2} alkyl, or $(CH_2)_{2-4}NR_AR_B$; (l) R_1 is H or (C_{3-4} cycloalkyl) C_{1-2} alkyl; (m) R_2 is H, methyl, C_{2-4} alkynyl, C_{3-5} cycloalkyl, or (C_{3-5} cycloalkyl)methyl; (n) R_A is H,
 20 methyl, ethyl, isobutyl, hydroxyethyl, hydroxypropyl, cyclopropylmethyl, cyclobutylmethyl, C_{3-4} alkynyl, phenyl, 2-piperidin-1-yl-ethyl, 2,3-dihydroxypropyl, 3-[4-(2-hydroxyethyl)-piperazin-1-yl]-propyl, 2-pyrrolidin-1-yl-ethyl, or 2-diethylamino-ethyl; and R_B is H; or where R_B is methyl and R_A is phenyl; (o) R_7 is $(CH_2)_{2-4}(NR_CR_D)$; (p) NR_CR_D is selected from morpholinyl, piperazinyl, pyrrolidinyl,
 25 or piperadiny; (q) R_C is methyl, ethyl, hydroxyethyl, or hydroxypropyl; (r) R_5 is methyl or chloro; (s) R_D is methyl, ethyl, hydroxyethyl, or hydroxypropyl; (t) or combinations thereof, such as wherein each of R_C and R_D is methyl or ethyl.

Preferably, where one of R_1 , R_2 , R_A , R_B , R_C , or R_D is an alkenyl or alkynyl group, the double or triple bond, respectively, is not adjacent the point of
 30 attachment. For example, where W is NR_2OR_1 , R_2 is preferably prop-2-ynyl, or but-2 or 3-enyl, and less preferably prop-1-ynyl or but-1-enyl.

Examples of compounds of formula (I) include: 2-(2-chloro-4-iodo-phenylamino)-4-sulfamoyl-benzoic acid; 2-(2-chloro-4-iodo-phenylamino)-N-hydroxy-4-sulfamoyl-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-4-sulfamoyl-benzamide; 2-(2-chloro-4-iodo-phenylamino)-4-(2-morpholin-4-yl-ethylsulfamoyl)-benzoic acid; 2-(2-chloro-4-iodo-phenylamino)-N-hydroxy-4-(2-morpholin-4-yl-ethylsulfamoyl)-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-4-(2-morpholin-4-yl-ethylsulfamoyl)-benzamide; 2-(2-chloro-4-iodo-phenylamino)-3,4-difluoro-5-sulfamoyl-benzoic acid; 2-(2-chloro-4-iodo-phenylamino)-3,4-difluoro-N-hydroxy-5-sulfamoyl-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-sulfamoyl-benzamide; 2-(2-chloro-4-iodo-phenylamino)-3,4-difluoro-5-(2-morpholin-4-yl-ethylsulfamoyl)-benzoic acid; 2-(2-chloro-4-iodo-phenylamino)-3,4-difluoro-N-hydroxy-5-(2-morpholin-4-yl-ethylsulfamoyl)-benzamide; and 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-(2-morpholin-4-yl-ethylsulfamoyl)-benzamide.

Other examples include 5-[bis-(4-methoxy-benzyl)-sulfamoyl]-2-(2-chloro-4-iodo-phenylamino)-3,4-difluoro-benzoic acid; and 2-(2-chloro-4-iodo-phenylamino)-5-dimethylsulfamoyl-3,4-difluoro-benzoic acid methyl ester.

Additional examples include 5-(bis-pyridin-3-ylmethyl-sulfamoyl)-3,4-difluoro-2-(4-iodo-phenylamino)-benzoic acid; 5-(bis-pyridin-3-ylmethyl-sulfamoyl)-N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-phenylamino)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-phenylamino)-5-(methyl-pyridin-3-ylmethyl-sulfamoyl)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-phenylamino)-5-[(pyridin-3-ylmethyl)-sulfamoyl]-benzamide; N-cyclopropylmethoxy-5-[(3-diethylamino-propyl)-pyridin-3-ylmethyl-sulfamoyl]-3,4-difluoro-2-(4-iodo-phenylamino)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-5-[(3-hydroxy-propyl)-pyridin-3-ylmethyl-sulfamoyl]-2-(4-iodo-phenylamino)-benzamide; N-cyclopropylmethoxy-5-(ethyl-pyridin-3-ylmethyl-sulfamoyl)-3,4-difluoro-2-(4-iodo-phenylamino)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-5-[(2-hydroxy-ethyl)-pyridin-3-ylmethyl-sulfamoyl]-2-(4-iodo-phenylamino)-benzamide; 5-(bis-pyridin-2-ylmethyl-sulfamoyl)-3,4-difluoro-2-(4-iodo-phenylamino)-benzoic acid; 5-(bis-pyridin-2-ylmethyl-sulfamoyl)-N-

cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-phenylamino)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-phenylamino)-5-(methyl-pyridin-2-ylmethyl-sulfamoyl)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-phenylamino)-5-[(pyridin-2-ylmethyl)-sulfamoyl]-benzamide; 5-(bis-pyridin-3-ylmethyl-sulfamoyl)-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzoic acid; 5-(bis-pyridin-3-ylmethyl-sulfamoyl)-N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-5-(methyl-pyridin-3-ylmethyl-sulfamoyl)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-5-[(pyridin-3-ylmethyl)-sulfamoyl]-benzamide; N-cyclopropylmethoxy-5-[(3-diethylamino-propyl)-pyridin-3-ylmethyl-sulfamoyl]-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-5-[(3-hydroxy-propyl)-pyridin-3-ylmethyl-sulfamoyl]-2-(4-iodo-2-methyl-phenylamino)-benzamide; N-cyclopropylmethoxy-5-(ethyl-pyridin-3-ylmethyl-sulfamoyl)-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-5-[(2-hydroxy-ethyl)-pyridin-3-ylmethyl-sulfamoyl]-2-(4-iodo-2-methyl-phenylamino)-benzamide; 5-(bis-pyridin-2-ylmethyl-sulfamoyl)-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzoic acid; 5-(bis-pyridin-2-ylmethyl-sulfamoyl)-N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-5-(methyl-pyridin-2-ylmethyl-sulfamoyl)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-5-[(pyridin-2-ylmethyl)-sulfamoyl]-benzamide; 5-(bis-pyridin-3-ylmethyl-sulfamoyl)-2-(2-chloro-4-iodo-phenylamino)-3,4-difluoro-benzoic acid, 5-(bis-pyridin-3-ylmethyl-sulfamoyl)-2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-(methyl-pyridin-3-ylmethyl-sulfamoyl)-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-[(pyridin-3-ylmethyl)-sulfamoyl]-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-5-[(3-diethylamino-propyl)-pyridin-3-ylmethyl-sulfamoyl]-3,4-difluoro-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-[(3-hydroxy-propyl)-pyridin-3-ylmethyl-sulfamoyl]-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-5-

- (ethyl-pyridin-3-ylmethyl-sulfamoyl)-3,4-difluoro-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-[(2-hydroxy-ethyl)-pyridin-3-ylmethyl-sulfamoyl]-benzamide; 5-(bis-pyridin-2-ylmethyl-sulfamoyl)-2-(2-chloro-4-iodo-phenylamino)-3,4-difluoro-benzoic acid; 5-(bis-pyridin-2-ylmethyl-
- 5 sulfamoyl)-2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-(methyl-pyridin-2-ylmethyl-sulfamoyl)-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-[(pyridin-2-ylmethyl)-sulfamoyl]-benzamide; N-cyclopropylmethoxy-3,4-difluoro-5-[(3-hydroxy-propyl)-pyridin-2-ylmethyl-sulfamoyl]-2-(4-iodo-phenylamino)-benzamide; N-
- 10 cyclopropylmethoxy-3,4-difluoro-5-[(2-hydroxy-ethyl)-pyridin-2-ylmethyl-sulfamoyl]-2-(4-iodo-phenylamino)-benzamide; 5-(benzyl-pyridin-2-ylmethyl-sulfamoyl)-N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-phenylamino)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-phenylamino)-5-[(pyridin-4-ylmethyl)-sulfamoyl]-benzamide; N-cyclopropylmethoxy-5-(ethyl-
- 15 pyridin-4-ylmethyl-sulfamoyl)-3,4-difluoro-2-(4-iodo-phenylamino)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-phenylamino)-5-(methyl-pyridin-4-ylmethyl-sulfamoyl)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-5-[(3-hydroxy-propyl)-pyridin-4-ylmethyl-sulfamoyl]-2-(4-iodo-phenylamino)-benzamide; N-
- 20 cyclopropylmethoxy-3,4-difluoro-5-[(2-hydroxy-ethyl)-pyridin-4-ylmethyl-sulfamoyl]-2-(4-iodo-phenylamino)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-phenylamino)-5-(methyl-phenyl-sulfamoyl)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-phenylamino)-5-phenylsulfamoyl-benzamide; N-Cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-phenylamino)-5-
- 25 (pyridin-3-ylsulfamoyl)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-5-[(3-hydroxy-propyl)-pyridin-2-ylmethyl-sulfamoyl]-2-(4-iodo-2-methyl-phenylamino)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-5-[(2-hydroxy-ethyl)-pyridin-2-ylmethyl-sulfamoyl]-2-(4-iodo-2-methyl-phenylamino)-benzamide; 5-(benzyl-pyridin-2-ylmethyl-sulfamoyl)-N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-2-
- 30 methyl-phenylamino)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-5-[(pyridin-4-ylmethyl)-sulfamoyl]-benzamide; N-cyclopropylmethoxy-5-(ethyl-pyridin-4-ylmethyl-sulfamoyl)-3,4-difluoro-2-(4-iodo-

2-methyl-phenylamino)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-5-(methyl-pyridin-4-ylmethyl-sulfamoyl)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-5-[(3-hydroxy-propyl)-pyridin-4-ylmethyl-sulfamoyl]-2-(4-iodo-2-methyl-phenylamino)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-5-[(2-hydroxy-ethyl)-pyridin-4-ylmethyl-sulfamoyl]-2-(4-iodo-2-methyl-phenylamino)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-5-(methyl-phenyl-sulfamoyl)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-5-phenylsulfamoyl-benzamide; N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-5-(pyridin-3-ylsulfamoyl)-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-[(3-hydroxy-propyl)-pyridin-2-ylmethyl-sulfamoyl]-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-[(2-hydroxy-ethyl)-pyridin-2-ylmethyl-sulfamoyl]-benzamide; 5-(benzyl-pyridin-2-ylmethyl-sulfamoyl)-2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-[(pyridin-4-ylmethyl)-sulfamoyl]-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-5-(ethyl-pyridin-4-ylmethyl-sulfamoyl)-3,4-difluoro-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-(methyl-pyridin-4-ylmethyl-sulfamoyl)-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-[(3-hydroxy-propyl)-pyridin-4-ylmethyl-sulfamoyl]-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-[(2-hydroxy-ethyl)-pyridin-4-ylmethyl-sulfamoyl]-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-(methyl-phenyl-sulfamoyl)-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-phenylsulfamoyl-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-(pyridin-3-ylsulfamoyl)-benzamide; N-cyclopropylmethoxy-2-(4-iodo-phenylamino)-4-phenylsulfamoyl-benzamide; N-cyclopropylmethoxy-2-(4-iodo-phenylamino)-4-(pyridin-3-ylsulfamoyl)-benzamide; N-cyclopropylmethoxy-2-(4-iodo-phenylamino)-4-[(pyridin-3-ylmethyl)-sulfamoyl]-benzamide; 4-(bis-pyridin-3-ylmethyl-sulfamoyl)-N-cyclopropylmethoxy-2-(4-iodo-phenylamino)-benzamide; N-cyclopropylmethoxy-4-[(2-hydroxy-ethyl)-pyridin-4-ylmethyl-sulfamoyl]-2-(4-

iodo-phenylamino)-benzamide; N-cyclopropylmethoxy-2-(4-iodo-phenylamino)-4-
 (methyl-pyridin-3-ylmethyl-sulfamoyl)-benzamide; N-cyclopropylmethoxy-4-[(3-
 diethylamino-propyl)-pyridin-3-ylmethyl-sulfamoyl]-2-(4-iodo-phenylamino)-
 benzamide; N-cyclopropylmethoxy-2-(4-iodo-2-methyl-phenylamino)-4-
 5 phenylsulfamoyl-benzamide; N-cyclopropylmethoxy-2-(4-iodo-2-methyl-
 phenylamino)-4-(pyridin-3-ylsulfamoyl)-benzamide; N-cyclopropylmethoxy-2-(4-
 iodo-2-methyl-phenylamino)-4-[(pyridin-3-ylmethyl)-sulfamoyl]-benzamide; 4-(bis-
 pyridin-3-ylmethyl-sulfamoyl)-N-cyclopropylmethoxy-2-(4-iodo-2-methyl-
 phenylamino)-benzamide; N-cyclopropylmethoxy-4-[(2-hydroxy-ethyl)-pyridin-4-
 10 ylmethyl-sulfamoyl]-2-(4-iodo-2-methyl-phenylamino)-benzamide; N-
 cyclopropylmethoxy-2-(4-iodo-2-methyl-phenylamino)-4-(methyl-pyridin-3-
 ylmethyl-sulfamoyl)-benzamide; N-cyclopropylmethoxy-4-[(3-diethylamino-
 propyl)-pyridin-3-ylmethyl-sulfamoyl]-2-(4-iodo-2-methyl-phenylamino)-
 benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-4-
 15 phenylsulfamoyl-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-
 cyclopropylmethoxy-4-(pyridin-3-ylsulfamoyl)-benzamide; 2-(2-chloro-4-iodo-
 phenylamino)-N-cyclopropylmethoxy-4-[(pyridin-3-ylmethyl)-sulfamoyl]-
 benzamide; 4-(bis-pyridin-3-ylmethyl-sulfamoyl)-2-(2-chloro-4-iodo-phenylamino)-
 N-cyclopropylmethoxy-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-
 20 cyclopropylmethoxy-4-[(2-hydroxy-ethyl)-pyridin-4-ylmethyl-sulfamoyl]-
 benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-4-(methyl-
 pyridin-3-ylmethyl-sulfamoyl)-benzamide; and 2-(2-chloro-4-iodo-phenylamino)-N-
 cyclopropylmethoxy-4-[(3-diethylamino-propyl)-pyridin-3-ylmethyl-sulfamoyl]-
 benzamide.

25 Further examples include : PD 298469, 2-(2-Chloro-4-iodo-phenylamino)-
 3,4-difluoro-N-methoxy-5-(4-methyl-piperazine-1-sulfonyl)-benzamide; PD
 298470, 2-(2-Chloro-4-iodo-phenylamino)-5-[(2-diethylamino-ethyl)-methyl-
 sulfamoyl]-3,4-difluoro-N-methoxy-benzamide; PD 298450, 2-(2-Chloro-4-iodo-
 phenylamino)-3,4-difluoro-N-methoxy-5-(methyl-prop-2-ynyl-sulfamoyl)-
 30 benzamide; PD 298451, 1-[4-(2-Chloro-4-iodo-phenylamino)-2,3-difluoro-5-
 methoxycarbamoyl-benzenesulfonyl]-piperidine-3-carboxylic acid amide;


PD 298452, 2-(2-Chloro-4-iodo-phenylamino)-3,4-difluoro-N-methoxy-5-[methyl-(2-pyridin-2-yl-ethyl)-sulfamoyl]-benzamide; PD 298453, 2-(2-Chloro-4-iodo-phenylamino)-5-[(3-dimethylamino-propyl)-methyl-sulfamoyl]-3,4-difluoro-N-methoxy-benzamide; PD 298454, 2-(2-Chloro-4-iodo-phenylamino)-3,4-difluoro-
5 N-methoxy-5-(4-pyridin-2-yl-piperazine-1-sulfonyl)-benzamide; PD 298455, 5-[Bis-(2-methoxy-ethyl)-sulfamoyl]-2-(2-chloro-4-iodo-phenylamino)-3,4-difluoro-N-methoxy-benzamide; PD 298456, 5-[Benzyl-(2-dimethylamino-ethyl)-sulfamoyl]-2-(2-chloro-4-iodo-phenylamino)-3,4-difluoro-N-methoxy-benzamide; and PD
298457, N-Allyloxy-2-(2-chloro-4-iodo-phenylamino)-5-dimethylsulfamoyl-3,4-
10 difluoro-benzamide; PD 298461, N-Allyloxy-2-(2-chloro-4-iodo-phenylamino)-3,4-difluoro-5-(methyl-prop-2-ynyl-sulfamoyl)-benzamide; PD 298462, N-Allyloxy-2-(2-chloro-4-iodo-phenylamino)-3,4-difluoro-5-[4-(4-fluoro-phenyl)-piperazine-1-sulfonyl]-benzamide; PD 298466, N-Allyloxy-5-[benzyl-(2-dimethylamino-ethyl)-sulfamoyl]-2-(2-chloro-4-iodo-phenylamino)-3,4-difluoro-benzamide; PD 298468,
15 2-(2-Chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-(4-methyl-piperazine-1-sulfonyl)-benzamide; and PD 298449, 2-(2-Chloro-4-iodo-phenylamino)-3,4-difluoro-5-(methoxy-methyl-sulfamoyl)-N-(2-morpholin-4-yl-ethoxy)-benzamide.

Particularly preferred compounds include: PD 298458, N-Allyloxy-2-(2-
20 chloro-4-iodo-phenylamino)-3,4-difluoro-5-(4-methyl-piperazine-1-sulfonyl)-benzamide; PD 298459, N-Allyloxy-2-(2-chloro-4-iodo-phenylamino)-3,4-difluoro-5-(methyl-phenyl-sulfamoyl)-benzamide; PD 298460, 5-(Allyl-methyl-sulfamoyl)-N-allyloxy-2-(2-chloro-4-iodo-phenylamino)-3,4-difluoro-benzamide; PD 298463,
1-[5-Allyloxycarbamoyl-4-(2-chloro-4-iodo-phenylamino)-2,3-difluoro-
25 benzenesulfonyl]-piperidine-3-carboxylic acid amide; PD 298464, N-Allyloxy-2-(2-chloro-4-iodo-phenylamino)-5-[(3-dimethylamino-propyl)-methyl-sulfamoyl]-3,4-difluoro-benzamide; PD 298465, N-Allyloxy-2-(2-chloro-4-iodo-phenylamino)-3,4-difluoro-5-(4-pyridin-2-yl-piperazine-1-sulfonyl)-benzamide; and PD 298467, N-Allyloxy-2-(2-chloro-4-iodo-phenylamino)-3,4-difluoro-5-(methoxy-methyl-
30 sulfamoyl)-benzamide.


C. Synthesis

The disclosed compounds can be synthesized according to the following four Schemes, or variants thereof. These synthetic strategies, which are suitable
5 for conventional or combinatorial synthetic methods, are further exemplified in Examples 1-4 below.


Scheme 1




Scheme 2



Scheme 3



Scheme 4



- Amine reagents such as R_6R_7NH in the schemes above are either commercially available or through straightforward modification of commercially available intermediates. Examples of such amine reagents, which can be reacted with the appropriate intermediate in a combinatorial or matrix method, are
- 5 provided below. For example, in section B (Compounds), starting at page 8, line 16, there are three sets of thirty (one set each for $R_5 = H, Me, \text{ and } Cl$). The table below provides a number (corresponding to order that the name is found in the text; for example, "1" corresponds to compounds 1, 31, and 61 in the list of 90 compounds); the amine reagent name; and a Chemical Abstracts number.
- 10 Where a PD number is listed, the amine reagent was prepared from commercially available starting materials.

Number (position in subset of 30)	Amine reagent name	CAS # or PD #
1	3,3'-dipicolylamine	1656-94-6
2	3,3'-dipicolylamine	1656-94-6
3	3-(methylaminomethyl)pyridine	20173-04-0
4	3-(aminomethyl)pyridine	3731-52-0
5	"N-(3-diethylaminopropyl)-N-(pyridin-3-ylmethyl)amine"	PD 0096419
6	3-(3-pyridylmethylamino)-1-propanol	6951-00-4
7	3-(ethylaminomethyl)pyridine	PD 0133573
8	2-(3-pyridylmethylamino)ethanol hydrochloride	PD 0018185-0002
9	di-(2-picolyl)amine	1539-42-0
10	di-(2-picolyl)amine	1539-42-0
11	2-(methylaminomethyl)pyridine	PD 0091430
12	2-(aminomethyl)pyridine	3731-51-9
13	3-(2-pyridylmethylamino)-1-propanol	6950-99-8
14	2-(2-pyridylmethylamino)ethanol	PD 0018354
15	2-(N-benzylaminomethyl)pyridine	PD 0054372
16	4-(aminomethyl)pyridine	3731-53-1
17	4-(ethylaminomethyl)pyridine	33403-97-3
18	4-(methylaminomethyl)pyridine	PD 0111199
19	3-(4-pyridylmethylamino)-1-propanol	7251-62-9
20	2-(4-pyridylmethylamino)ethanol hydrochloride	PD 0018008-0002
21	N-methylaniline	100-61-8
22	aniline	62-53-3

23	3-aminopyridine	462-08-8
24	<i>aniline</i>	62-53-3
25	3-aminopyridine	462-08-8
26	3-(aminomethyl)pyridine	3731-52-0
27	3,3'-dipicolylamine	1656-94-6
28	2-(4-pyridylmethylamino)ethanol hydrochloride	<u>PD 0018008-0002</u>
29	3-(methyaminomethyl)pyridine	20173-04-0
30	"N-(3-diethylaminopropyl)-N-(pyridin-3-ylmethyl)amine"	<u>PD 0096419</u>

Additional compounds within claim 1 can be made with the following amine reagents. The corresponding CAS number is provided.

5	2-(methylamino)pyridine	4597-87-9
	2-benzylaminopyridine	6935-27-9
	2-allylaminopyridine	5866-28-4
	2,2'-dipyridylamine	1202-34-2
10	2-anilinopyridine	6631-37-4
	2-aminopyridine	504-29-0
	4-aminopyridine	504-24-5
	2-benzylaminopyridine	6935-27-9
	2-(4-methoxybenzyl)aminopyridine	52818-63-0
15	2-methylaminopyridine	4597-87-9

Combinatorial Synthesis

The following stock solutions were prepared:

- 20 1) An acetonitrile (anhydrous) stock solution 0.05 M in 5-chlorosulfonyl-2,3,4-trifluoro-benzoyl chloride.
- 2) Acetonitrile (anhydrous) stock solutions 0.05 M in each of the four appropriate hydroxylamine hydrochlorides (see list A) and 0.3 M in 2,6-lutidine.
- 3) Acetonitrile stock solutions 0.05 M in each of the 25 appropriate amines (see
- 25 list B). Note that amine salts that were not soluble were also 0.1 M in 2,6-lutidine.
- 4) Acetonitrile (anhydrous) stock solutions in each of the 3 appropriate anilines (see list C) and 0.88 M in lithium bis(trimethylsilyl)amide (1.0 M in tetrahydrofuran).

An array which treated 4 hydroxylamine hydrochlorides independently with 5-chlorosulfonyl-2,3,4-trifluoro-benzoyl chloride, 25 amines, and 1 aniline was prepared to yield a total of 100 reactions. A liquid handling robot was used to transfer the reagents in such a manner as to insure that all possible combinations were achieved. The appropriate hydroxylamine hydrochloride solution (0.05 mmol, 1 mL) was added to a 2-dram vial, and each vial was treated with 5-chlorosulfonyl-2,3,4-trifluoro-benzoyl chloride solution (0.05 mmol, 1 mL). After 20 minutes the appropriate amine solution (0.05 mmol, 1 mL) was added sequentially. After a further 20 minutes the vials were treated with the solution of 4-iodoaniline (0.055 mmol, 1 mL). The vials were capped and shaken overnight at room temperature. The reactions were quenched with 1 mL of a 1 M aqueous ammonium chloride solution. The vials were concentrated to dryness under a stream of nitrogen and purified by reverse phase HPLC using a 30x100 mm YMC ODS-A (C18) column. The mobile phase was acetonitrile/water (both with 0.05% trifluoroacetic acid) at 25 mL/min and a linear gradient of 10-100% over 6.5 min and then 3.5 min at 100%, detection was at 214 nm.

An array which treated 4 hydroxylamine hydrochlorides independently with 5-chlorosulfonyl-2,3,4-trifluoro-benzoyl chloride, 25 amines, and 1 aniline was prepared to yield a total of 100 reactions. A liquid handling robot was used to transfer the reagents in such a manner as to insure that all possible combinations were achieved. The appropriate hydroxylamine hydrochloride solution (0.05 mmol, 1 mL) was added to a 2-dram vial, and each vial was treated with 5-chlorosulfonyl-2,3,4-trifluoro-benzoyl chloride solution (0.05 mmol, 1 mL). After 20 minutes the appropriate amine solution (0.05 mmol, 1 mL) was added sequentially. After a further 20 minutes the vials were treated with the solution of 4-iodo-2-methylaniline (0.05 mmol, 0.91 mL). The vials were capped and shaken overnight at room temperature. The reactions were quenched with 1 mL of a 1 M aqueous ammonium chloride solution. The vials were concentrated to dryness under a stream of nitrogen and purified by reverse phase HPLC using a 30x100 mm YMC ODS-A (C18) column. The mobile phase was acetonitrile/water (both

with 0.05% trifluoroacetic acid) at 25 mL/min and a linear gradient of 10-100% over 6.5 min and then 3.5 min at 100%, detection was at 214 nm.

An array which treated 4 hydroxylamine hydrochlorides independently with 5-chlorosulfonyl-2,3,4-trifluoro-benzoyl chloride, 25 amines, and 1 aniline was prepared to yield a total of 100 reactions. A liquid handling robot was used to transfer the reagents in such a manner as to insure that all possible combinations were achieved. The appropriate hydroxylamine hydrochloride solution (0.05 mmol, 1 mL) was added to a 2-dram vial, and each vial was treated with 5-chlorosulfonyl-2,3,4-trifluoro-benzoyl chloride solution (0.05 mmol, 1 mL). After 20 minutes the appropriate amine solution (0.05 mmol, 1 mL) was added sequentially. After a further 20 minutes the vials were treated with the solution of 2-chloro-4-iodoaniline (0.05 mmol, 0.91 mL). The vials were capped and shaken overnight at room temperature. The reactions were quenched with 1 mL of a 1 M aqueous ammonium chloride solution. The vials were concentrated to dryness under a stream of nitrogen and purified by reverse phase HPLC using a 30x100 mm YMC ODS-A (C18) column. The mobile phase was acetonitrile/water (both with 0.05% trifluoroacetic acid) at 25 mL/min and a linear gradient of 10-100% over 6.5 min and then 3.5 min at 100%, detection was at 214 nm.

Combinatorial Synthesis
Table of Example Reagents

5 List A-Hydroxylamines:

1. O-methyl-hydroxylamine
2. O-allyl-hydroxylamine hydrochloride monohydrate (Aldrich)
3. O-cyclopropylmethyl-hydroxylamine hydrochloride
- 10 4. O-(2-morpholin-4-yl-ethyl)-hydroxylamine hydrochloride


List B-Amines:

1. dimethylamine
- 15 2. diethylamine
3. isopropyl-methyl-amine
4. diisopropylamine
5. methylhydrazine
6. 1-methylpiperazine
- 20 7. *N,N*-diethyl-*N'*-methylethane-1,2-diamine
8. benzylmethylamine
9. dibenzylamine
10. methyl-phenyl-amine
11. allyl-methyl-amine
- 25 12. methyl-prop-2-ynyl-amine
13. methylamino-acetonitrile hydrochloride
14. 1-(4-fluoro-phenyl)-piperazine
15. furan-2-ylmethyl-methyl-amine
16. piperidine-3-carboxylic acid amide
- 30 17. methyl-phenethyl-amine
18. methyl-(2-pyridin-2-yl-ethyl)-amine
19. *N,N,N'*-trimethyl-propane-1,3-diamine
20. methyl-(1-methyl-piperidin-4-yl)-amine
21. 1-pyridin-2-yl-piperazine
- 35 22. bis-(2-methoxy-ethyl)-amine
23. *N'*-benzyl-*N,N*-dimethyl-ethane-1,2-diamine
24. methylamino-acetic acid *tert*-butyl ester hydrochloride
25. *O,N*-dimethyl-hydroxylamine hydrochloride

40

List C-Anilines:

1. 4-iodoaniline
- 45 2. 2-chloro-4-iodoaniline
3. 4-iodo-2-methylaniline



D. Uses

The disclosed compositions are useful as both prophylactic and therapeutic treatments for diseases or conditions as provided in the Summary section, as well as diseases or conditions modulated by the MEK cascade. Examples include stroke, heart failure, osteoarthritis, rheumatoid arthritis, organ transplant rejection, and a variety of tumors such as ovarian, lung, pancreatic, brain, prostatic, renal, and colorectal.

1. Dosages

Those skilled in the art will be able to determine, according to known methods, the appropriate dosage for a patient, taking into account factors such as age, weight, general health, the type of symptoms requiring treatment, and the presence of other medications. In general, an effective amount will be between 0.1 and 1000 mg/kg per day, preferably between 1 and 300 mg/kg body weight, and daily dosages will be between 10 and 5000 mg for an adult subject of normal weight. Capsules, tablets or other formulations (such as liquids and film-coated tablets) may be of between 5 and 200 mg, such as 10, 15, 25, 35, 50 mg, 60 mg, and 100 mg and can be administered according to the disclosed methods.

2. Formulations

Dosage unit forms include tablets, capsules, pills, powders, granules, aqueous and nonaqueous oral solutions and suspensions, and parenteral solutions packaged in containers adapted for subdivision into individual doses. Dosage unit forms can also be adapted for various methods of administration, including controlled release formulations, such as subcutaneous implants. Administration methods include oral, rectal, parenteral (intravenous, intramuscular, subcutaneous), intracisternal, intravaginal, intraperitoneal, intravesical, local (drops, powders, ointments, gels, or cream), and by inhalation (a buccal or nasal spray).

Parenteral formulations include pharmaceutically acceptable aqueous or nonaqueous solutions, dispersion, suspensions, emulsions, and sterile powders for the preparation thereof. Examples of carriers include water, ethanol, polyols (propylene glycol, polyethylene glycol), vegetable oils, and injectable organic

esters such as ethyl oleate. Fluidity can be maintained by the use of a coating such as lecithin, a surfactant, or maintaining appropriate particle size. Carriers for solid dosage forms include (a) fillers or extenders, (b) binders, (c) humectants, (d) disintegrating agents, (e) solution retarders, (f) absorption
5 accelerators, (g) adsorbants, (h) lubricants, (i) buffering agents, and (j) propellants.

Compositions may also contain adjuvants such as preserving, wetting, emulsifying, and dispensing agents; antimicrobial agents such as parabens, chlorobutanol, phenol, and sorbic acid; isotonic agents such as a sugar or sodium
10 chloride; absorption-prolonging agents such as aluminum monostearate and gelatin; and absorption-enhancing agents.

3. Related compounds

The invention provides the disclosed compounds and closely related, pharmaceutically acceptable forms of the disclosed compounds, such as salts,
15 esters, amides, hydrates or solvated forms thereof; masked or protected forms; and racemic mixtures, or enantiomerically or optically pure forms.

Pharmaceutically acceptable salts, esters, and amides include carboxylate salts (e.g., C₁₋₈ alkyl, cycloalkyl, aryl, heteroaryl, or non-aromatic heterocyclic), amino acid addition salts, esters, and amides which are within a reasonable
20 benefit/risk ratio, pharmacologically effective, and suitable for contact with the tissues of patients without undue toxicity, irritation, or allergic response. Representative salts include hydrobromide, hydrochloride, sulfate, bisulfate, nitrate, acetate, oxalate, valerate, oleate, palmitate, stearate, laurate, borate, benzoate, lactate, phosphate, tosylate, citrate, maleate, fumarate, succinate,
25 tartrate, naphthylate, mesylate, glucoheptonate, lactobionate, and laurylsulfonate. These may include alkali metal and alkali earth cations such as sodium, potassium, calcium, and magnesium, as well as non-toxic ammonium, quaternary ammonium, and amine cations such as tetramethyl ammonium, methylamine, trimethylamine, and ethylamine. See, for example, S.M. Berge, et al., "Pharmaceutical Salts," *J. Pharm. Sci.*, 1977, 66:1-19 which is incorporated
30 herein by reference. Representative pharmaceutically acceptable amides of the invention include those derived from ammonia, primary C₁₋₆ alkyl amines and

secondary di (C₁₋₆ alkyl) amines. Secondary amines include 5- or 6-membered heterocyclic or heteroaromatic ring moieties containing at least one nitrogen atom and optionally between 1 and 2 additional heteroatoms. Preferred amides are derived from ammonia, C₁₋₃ alkyl primary amines, and di (C₁₋₂ alkyl)amines.

- 5 Representative pharmaceutically acceptable esters of the invention include C₁₋₇ alkyl, C₅₋₇ cycloalkyl, phenyl, and phenyl(C₁₋₆)alkyl esters. Preferred esters include methyl esters.

- The invention also includes disclosed compounds having one or more functional groups (e.g., hydroxyl, amino, or carboxyl) masked by a protecting
10 group. Some of these masked or protected compounds are pharmaceutically acceptable; others will be useful as intermediates. Synthetic intermediates and processes disclosed herein, and minor modifications thereof, are also within the scope of the invention.

15 HYDROXYL PROTECTING GROUPS

Hydroxyl protecting groups include: ethers, esters, and protection for 1,2- and 1,3-diols. The ether protecting groups include: methyl, substituted methyl ethers, substituted ethyl ethers, substituted benzyl ethers, silyl ethers and conversion of silyl ethers to other functional groups.

20 Substituted Methyl Ethers

- Substituted methyl ethers include: methoxymethyl, methylthiomethyl, *t*-utylthiomethyl, (phenyldimethylsilyl) methoxymethyl, benzyloxymethyl, *p*-ethoxybenzyloxymethyl, (4-methoxyphenoxy) methyl, guaiacolmethyl, *t*-butoxymethyl, 4-pentenylloxymethyl, siloxymethyl, 2-methoxyethoxymethyl,
25 2,2,2-trichloroethoxymethyl, bis(2-chloro- ethoxy)methyl, 2-(trimethylsilyl)-ethoxymethyl, tetrahydropyranyl, 3-bromotetrahydro-pyranyl, tetrahydrothiopyranyl, 1-methoxycyclohexyl, 4-methoxytetrahydropyranyl, 4-methoxytetrahydro- thiopyranyl, 4-methoxytetrahydrothiopyranyl *S,S*-dioxido, 1-[(2-chloro-4-methyl)phenyl]-4-methoxypiperidin-4-yl, 1,4-dioxan-2-yl,
30 tetrahydrofuranyl, tetrahydrothiofuranyl, and 2,3,3a,4,5,6,7,7a-octahydro-7,8,8-trimethyl-4,7-ethanobenzofuran-2-yl.

Substituted Ethyl Ethers

Substituted ethyl ethers include: 1-ethoxyethyl, 1-(2-chloroethoxy)ethyl, 1-methyl-1-methoxyethyl, 1-methyl-1-benzyloxyethyl, 1-methyl-1-benzyloxy-2-fluoroethyl, 2,2,2-trichloroethyl, 2-trimethylsilyl, 2-(phenylselenyl)ethyl, *t*-butyl, allyl, *p*-chlorophenyl, *p*-methoxyphenyl, 2,4-dinitrophenyl, and benzyl.

Substituted Benzyl Ethers

Substituted benzyl ethers include: *p*-methoxybenzyl, 3,4-dimethoxybenzyl, *o*-nitrobenzyl, *p*-nitrobenzyl, *p*-halobenzyl, 2,6-dichlorobenzyl, *p*-cyanobenzyl, *p*-phenylbenzyl, 2- and 4-picolyl, 3-methyl-2-picolyl *N*-oxido, diphenylmethyl, *p*, *p*'-dinitrobenzhydryl, 5-dibenzosuberone, triphenylmethyl, α -naphthylidiphenylmethyl, *p*-methoxyphenyldiphenylmethyl, di(*p*-methoxyphenyl)phenylmethyl, tri(*p*-methoxyphenyl)methyl, 4-(4'-bromophenacyloxy)phenyldiphenylmethyl, 4,4',4''-tris(4,5-dichlorophthalimidophenyl)methyl, 4,4',4''-tris(levulinoyloxyphenyl)methyl, 4,4',4''-tris(benzoyloxyphenyl)methyl, 3-(imidazol-1-ylmethyl)bis(4',4''-dimethoxyphenyl)-methyl, 1,1-bis(4-methoxyphenyl)-1'-pyrenylmethyl, 9-anthryl, 9-(9-phenyl) xanthenyl, 9-(9-phenyl-10-oxo) anthryl, 1,3-benzodithiolan-2-yl, and benzoisothiazolyl *S,S*-dioxide.

Silyl Ethers

Silyl ethers include: trimethylsilyl, triethylsilyl, triisopropylsilyl, dimethylisopropylsilyl, diethylisopropylsilyl, dimethylhexylsilyl, *t*-butyldimethylsilyl, *t*-butyldiphenylsilyl, tribenzylsilyl, tri-*p*-xylylsilyl, triphenylsilyl, diphenylmethylsilyl, and *t*-butylmethoxy-phenylsilyl.

ESTERS

Esters protecting groups include: esters, carbonates, assisted cleavage, miscellaneous esters, and sulfonates.

Esters

Examples of protective esters include: formate, benzoylformate, acetate, chloroacetate, dichloroacetate, trichloroacetate, trifluoroacetate, methoxyacetate, triphenylmethoxyacetate, phenoxyacetate, *p*-chlorophenoxyacetate, *p*-phenylacetate, 3-phenylpropionate, 4-oxopentanoate (levulinate), 4,4-

(ethylenedithio) pentanoate, pivaloate, adamantate, crotonate, 4-methoxycrotonate, benzoate, *p*-phenylbenzoate, and 2,4,6-trimethylbenzoate (mesitoate).

Carbonates

- 5 Carbonates include: methyl, 9-fluorenylmethyl, ethyl, 2,2,2-trichloroethyl, 2-(trimethylsilyl) ethyl, 2-(phenylsulfonyl) ethyl, 2-(triphenylphosphonio) ethyl, isobutyl, vinyl, allyl, *p*-nitrophenyl, benzyl, *p*-methoxybenzyl, 3,4-dimethoxybenzyl, *o*-nitrobenzyl, *p*-nitrobenzyl, *S*-benzyl thiocarbonate, 4-ethoxy-1-naphthyl, and methyl dithiocarbonate.

10 Assisted Cleavage

Examples of assisted cleavage protecting groups include: 2-iodobenzoate, 4-azido-butylate, 4-nitro-4-methylpentanoate, *o*-(dibromomethyl) benzoate, 2-formylbenzene-sulfonate, 2-(methylthiomethoxy) ethyl carbonate, 4-(methylthiomethoxymethyl) benzoate, and 2-(methylthiomethoxymethyl)

- 15 benzoate.

Miscellaneous Esters

- In addition to the above classes, miscellaneous esters include: 2,6-dichloro-4-methylphenoxyacetate, 2,6-dichloro-4-(1,1,3,3-tetramethylbutyl) phenoxyacetate, 20 2,4-bis(1,1-dimethylpropyl) phenoxyacetate, chlorodiphenylacetate, isobutyrate, monosuccinoate, (*E*)-2-methyl-2-butenate (tigloate), *o*-(methoxycarbonyl) benzoate, *p*-P-benzoate, α -naphthoate, nitrate, alkyl *N,N,N',N'*-tetramethylphosphorodiamidate, *N*-phenylcarbamate, borate, dimethylphosphinothioyl, and 2,4-dinitrophenylsulfenate.

25 Sulfonates

Protective sulfates includes: sulfate, methanesulfonate(mesylate), benzylsulfonate, and tosylate.

PROTECTION FOR 1,2- AND 1,3-DIOLS

- 30 The protection for 1,2 and 1,3-diols group includes: cyclic acetals and ketals, cyclic ortho esters, and silyl derivatives.

Cyclic Acetals and Ketals

- Cyclic acetals and ketals include: methylene, ethylidene, 1-*t*-butylethylidene, 1-phenylethylidene, (4-methoxyphenyl) ethylidene, 2,2,2-trichloroethylidene, acetonide (isopropylidene), cyclopentylidene, cyclohexylidene, cycloheptylidene,
- 5 benzylidene, *p*-methoxybenzylidene, 2,4-dimethoxybenzylidene, 3,4-dimethoxybenzylidene, and 2-nitrobenzylidene.

Cyclic Ortho Esters

- Cyclic ortho esters include: methoxymethylene, ethoxymethylene, dimethoxymethylene, 1-methoxyethylidene, 1-ethoxyethylidene, 1,2-dimethoxyethylidene,
- 10 α -methoxybenzylidene, 1-(*N,N*-dimethylamino)ethylidene derivative, α -(*N,N*-dimethylamino) benzylidene derivative, and 2-oxacyclopentylidene.

PROTECTION FOR THE CARBOXYL GROUPESTERS

- 15 Ester protecting groups include: esters, substituted methyl esters, 2-substituted ethyl esters, substituted benzyl esters, silyl esters, activated esters, miscellaneous derivatives, and stannyl esters.

Substituted Methyl Esters

- Substituted methyl esters include: 9-fluorenylmethyl, methoxymethyl,
- 20 methylthiomethyl, tetrahydropyranyl, tetrahydrofuranyl, methoxyethoxymethyl, 2-(trimethylsilyl)ethoxy-methyl, benzyloxymethyl, phenacyl, *p*-bromophenacyl, α -methylphenacyl, *p*-methoxyphenacyl, carboxamidomethyl, and *N*-phthalimidomethyl.

2-Substituted Ethyl Esters

- 25 2-Substituted ethyl esters include: 2,2,2-trichloroethyl, 2-haloethyl, 1-chloroalkyl, 2-(trimethylsilyl)ethyl, 2-methylthioethyl, 1,3-dithianyl-2-methyl, 2(*p*-nitrophenylsulfonyl)-ethyl, 2-(*p*-toluenesulfonyl)ethyl, 2-(2'-pyridyl)ethyl, 2-(diphenylphosphino)ethyl, 1-methyl-1-phenylethyl, *t*-butyl, cyclopentyl, cyclohexyl, allyl, 3-buten-1-yl, 4-(trimethylsilyl)-2-buten-1-yl, cinnamyl, α -methylcinnamyl,
- 30 phenyl, *p*-(methylmercapto)-phenyl, and benzyl.

Substituted Benzyl Esters

- Substituted benzyl esters include: triphenylmethyl, diphenylmethyl, bis(*o*-nitrophenyl)methyl, 9-anthrylmethyl, 2-(9,10-dioxo)anthrylmethyl, 5-dibenzo-suberyl, 1-pyrenylmethyl, 2-(trifluoromethyl)-6-chromylmethyl, 2,4,6-trimethylbenzyl, *p*-bromobenzyl, *o*-nitrobenzyl, *p*-nitrobenzyl, *p*-methoxybenzyl, 2,6-dimethoxybenzyl, 4-(methylsulfinyl)benzyl, 4-sulfobenzyl, piperonyl, and 4-*P*-benzyl.

Silyl Esters

- Silyl esters include: trimethylsilyl, triethylsilyl, *t*-butyldimethylsilyl, *i*-propyldimethylsilyl, phenyldimethylsilyl, and di-*t*-butylmethylsilyl.

Miscellaneous Derivatives

- Miscellaneous derivatives includes: oxazoles, 2-alkyl-1,3-oxazolines, 4-alkyl-5-oxo-1,3-oxazolidines, 5-alkyl-4-oxo-1,3-dioxolanes, ortho esters, phenyl group, and pentaaminocobalt(III) complex.

Stannyl Esters

- Examples of stannyl esters include: triethylstannyl and tri-*n*-butylstannyl.

AMIDES AND HYDRAZIDES

- Amides include: *N,N*-dimethyl, pyrrolidinyl, piperidinyl, 5,6-dihydrophenanthridinyl, *o*-nitroanilides, *N*-7-nitroindolyl, *N*-8-nitro-1,2,3,4-tetrahydroquinolyl, and *p*-*P*-benzenesulfonamides. Hydrazides include: *N*-phenyl, *N,N'*-diisopropyl and other dialkyl hydrazides.

PROTECTION FOR THE AMINO GROUP

25

CARBAMATES

Carbamates include: carbamates, substituted ethyl, assisted cleavage, photolytic cleavage, urea-type derivatives, and miscellaneous carbamates.

Carbamates

- Carbamates include: methyl and ethyl, 9-fluorenylmethyl, 9-(2-sulfo)fluorenylmethyl, 9-(2,7-dibromo)fluorenylmethyl, 2,7-di-*t*-butyl-[9-(10,10-dioxo-10,10,10,10-tetrahydro-thioxanthyl)]methyl, and 4-methoxyphenacyl.

Substituted Ethyl

Substituted ethyl protective groups include: 2,2,2-trichloroethyl, 2-trimethylsilylethyl, 2-phenylethyl, 1-(1-adamantyl)-1-methylethyl, 1,1-dimethyl-2-haloethyl, 1,1-dimethyl-2,2-dibromoethyl, 1,1-dimethyl-2,2,2-trichloroethyl, 1-methyl-1-(4-biphenyl)ethyl, 1-(3,5-di-*t*-butylphenyl)-1-methylethyl, 2-(2'-and 4'-pyridyl)ethyl, 2-(*N,N*-cyclohexylcarboxamido)-ethyl, *t*-butyl, 1-adamantyl, vinyl, allyl, 1-isopropylallyl, connamyl, 4-nitrocinnamyl, quinolyl, *N*-hydroxypiperidinyl, alkylidithio, benzyl, *p*-methoxybenzyl, *p*-nitrobenzyl, *p*-bromobenzyl, *p*-chlorobenzyl, 2,4-dichlorobenzyl, 4-methylsulfinylbenzyl, 9-anthrylmethyl, and diphenylmethyl.

Assisted Cleavage

Protection via assisted cleavage includes: 2-methylthioethyl, 2-methylsulfonylethyl, 2-(*p*-toluenesulfonyl)ethyl, [2-(1,3-dithianyl)]methyl, 4-methylthiophenyl, 2,4-dimethyl-thiophenyl, 2-phosphonioethyl, 2-triphenylphosphonioisopropyl, 1,1-dimethyl-2-cyanoethyl, *m*-chloro-*p*-acyloxybenzyl, *p*-(dihydroxyboryl)benzyl, 5-benzisoxazolyl-methyl, and 2-(trifluoromethyl)-6-chromonylmethyl.

Photolytic Cleavage

Photolytic cleavage methods use groups such as: *m*-nitrophenyl, 3,5-dimethoxybenzyl, *o*-nitrobenzyl, 3,4-dimethoxy-6-nitrobenzyl, and phenyl(*o*-nitrophenyl)methyl.

Urea-Type Derivatives

Examples of of urea-type derivatives include: phenothiazinyl-(10)-carbonyl derivative, *N*'-*p*-toluenesulfonylaminocarbonyl, and *N*'-phenylaminothiocarbonyl.

Miscellaneous Carbamates

In addition to the above, miscellaneous carbamates include: *t*-amyl, *S*-benzyl thiocarbamate, *p*-cyanobenzyl, cyclobutyl, cyclohexyl, cyclopentyl, cyclopropylmethyl, *p*-decyloxy-benzyl, diisopropylmethyl, 2,2-dimethoxycarbonylvinyl, *o*-(*N,N*-dimethyl-carboxamido)-benzyl, 1,1-dimethyl-3(*N,N*-dimethylcarboxamido)propyl, 1,1-dimethyl-propynyl, di(2-pyridyl)methyl, 2-

- furanylmethyl, 2-iodoethyl, isobornyl, isobutyl, isonicotinyl, *p*(*p*-methoxyphenylazo)benzyl, 1-methylcyclobutyl, 1-methylcyclohexyl, 1-methyl-1-cyclopropylmethyl, 1-methyl-(3,5-dimethoxyphenyl)ethyl, 1-methyl-1(*p*-henylazophenyl)-ethyl, 1-methyl-1-phenylethyl, 1-methyl-1-(4-pyridyl)ethyl, 5 phenyl, *p*-(phenylazo)benzyl, 2,4,6-tri-*t*-butylphenyl, 4-(trimethylammonium)-benzyl, and 2,4,6-trimethylbenzyl.

AMIDES

Amides

- 10 Amides includes: *N*-formyl, *N*-acetyl, *N*-chloroacetyl, *N*-trichloroacetyl, *N*-trifluoroacetyl, *N*-phenylacetyl, *N*-3-phenylpropionyl, *N*-picolinoyl, *N*-3-pyridyl-carboxamide, *N*-benzoylphenylalanyl derivative, *N*-benzoyl, and *N*-*p*-phenylbenzoyl.

15 Assisted Cleavage

- Assisted cleavage groups include: *N*-*o*-nitrophenylacetyl, *N*-*o*-nitrophenoxyacetyl, *N*-acetoacetyl, (*N*'-dithiobenzyloxycarbonylamino)acetyl, *N*-3-(*p*-hydroxyphenyl) propionyl, *N*-3-(*o*-nitrophenyl)propionyl, *N*-2-methyl-2-(*o*-nitrophenoxy)propionyl, *N*-2-methyl-2-(*o*-phenylazophenoxy)propionyl, *N*-4-20 chlorobutyryl, *N*-3-methyl-3-nitrobutyryl, *N*-*o*-nitrocinnamoyl, *N*-acetylmethionine derivative, *N*-*o*-nitrobenzoyl, *N*-*o*-(benzoyloxymethyl)benzoyl, and 4,5-diphenyl-3-oxazolin-2-one.

Cyclic Imide Derivatives

- Cyclic imide derivatives include: *N*-phthalimide, *N*-dithiasuccinoyl, 25 *N*-2,3-diphenyl-maleoyl, *N*-2,5-dimethylpyrrolyl, *N*-1,1,4,4-tetramethyldisilylazacyclopentane adduct, 5-substituted 1,3-dimethyl-1,3,5-triazacyclohexan-2-one, 5-substituted 1,3-dibenzyl-1,3,5-triazacyclohexan-2-one, and 1-substituted 3,5-dinitro-4-pyridonyl.

SPECIAL -NH PROTECTIVE GROUPS

- Protective groups for – NH include: *N*-alkyl and *N*-aryl amines, imine derivatives, enamine derivatives, and *N*-hetero atom derivatives (such as *N*-metal, *N*-N, *N*-P, *N*-Si, and *N*-S), *N*-sulfenyl, and *N*-sulfonyl.

N-Alkyl and *N*-Aryl Amines

- N*-alkyl and *N*-aryl amines include: *N*-methyl, *N*-allyl, *N*-[2-(trimethylsilyl)ethoxyl]-methyl, *N*-3-acetoxypentyl, *N*-(1-isopropyl-4-nitro-2-oxo-3-pyrrolin-3-yl), quaternary ammonium salts, *N*-benzyl, *N*-di(4-methoxyphenyl)methyl, *N*-5-dibenzosuberonyl, *N*-triphenylmethyl, *N*-(4-methoxyphenyl)diphenylmethyl, *N*-9-phenylfluorenyl, *N*-2,7-dichloro-9-fluorenylmethylene, *N*-ferrocenylmethyl, and *N*-2-picolyamine *N*'-oxide.

Imine Derivatives

- Imine derivatives include: *N*-1,1-dimethylthiomethylene, *N*-benzylidene, *N*-*p*-methoxybenzylidene, *N*-diphenylmethylene, *N*-[(2-pyridyl)mesityl]methylene, *N*-(*N*',*N*'-dimethylaminomethylene), *N*,*N*'-isopropylidene, *N*-*p*-nitrobenzylidene, *N*-salicylidene, *N*-5-chlorosalicylidene, *N*-(5-chloro-2-hydroxyphenyl)phenylmethylene, and *N*-cyclohexylidene.

Enamine Derivative

- An example of an enamine derivative is *N*-(5,5-dimethyl-3-oxo-1-cyclohexenyl).

N-Hetero Atom Derivatives

- N*-metal derivatives include: *N*-borane derivatives, *N*-diphenylborinic acid derivative, *N*-[phenyl(pentacarbonylchromium- or -tungsten)]carbenyl, and *N*-copper or *N*-zinc chelate. Examples of *N*-*N* derivatives include: *N*-nitro, *N*-nitroso, and *N*-oxide. Examples of *N*-P derivatives include: *N*-diphenylphosphinyl, *N*-dimethylthiophosphinyl, *N*-diphenylthiophosphinyl, *N*-dialkyl phosphoryl, *N*-dibenzyl phosphoryl, and *N*-diphenyl phosphoryl. Examples of *N*-sulfenyl derivatives include: *N*-benzenesulfenyl, *N*-*o*-nitrobenzenesulfenyl, *N*-2,4-dinitrobenzenesulfenyl, *N*-pentachlorobenzenesulfenyl, *N*-2-nitro-4-methoxy-benzenesulfenyl, *N*-triphenylmethylsulfenyl, and *N*-3-nitropyridinesulfenyl.

N-sulfonyl derivatives include: *N*-*p*-toluenesulfonyl, *N*-benzenesulfonyl, *N*-2,3,6-trimethyl-4-methoxybenzenesulfonyl, *N*-2,4,6-trimethoxybenzenesulfonyl, *N*-2,6-dimethyl-4-methoxy-benzenesulfonyl, *N*-pentamethylbenzenesulfonyl, *N*-2,3,5,6-tetramethyl-4-methoxybenzene-sulfonyl, *N*-4-methoxybenzenesulfonyl, 5 *N*-2,4,6-trimethylbenzenesulfonyl, *N*-2,6-dimethoxy-4-methylbenzenesulfonyl, *N*-2,2,5,7,8-pentamethylchroman-6-sulfonyl, *N*-methanesulfonyl, *N*- β -trimethylsilylethanesulfonyl, *N*-9-anthracenesulfonyl, *N*-4-(4',8'-dimethoxynaphthylmethyl)-benzenesulfonyl, *N*-benzylsulfonyl, *N*-trifluoromethylsulfonyl, and *N*-phenacylsulfonyl.

10

Disclosed compounds which are masked or protected may be prodrugs, compounds metabolized or otherwise transformed *in vivo* to yield a disclosed compound, e.g., transiently during metabolism. This transformation may be a hydrolysis or oxidation which results from contact with a bodily fluid such as 15 blood, or the action of acids, or liver, gastrointestinal, or other enzymes.

Features of the invention are further described in the examples below.

E. EXAMPLES

EXAMPLE 1

5

Preparation of 2-(2-chloro-4-iodo-phenylamino)-5-dimethylsulfamoyl-3,4-difluoro-benzoic acid methyl ester (APK IC₅₀=222 nM)

10

Step a: Preparation of 1-dimethylsulfamoyl-2,3,4-trifluorobenzene

To a gently stirring solution comprised of 2,3,4-trifluorobenzenesulfonyl chloride (5.70 g, 0.0247 mol) in 1,2-dichloroethane (200 ml) was introduced by bubbling gaseous anhydrous dimethylamine. The mixture became cloudy after several minutes and was subsequently washed with water (200 ml), 6 N aqueous hydrochloric acid (200 ml), brine (200 ml), was dried over anhydrous magnesium sulfate, and was concentrated *in vacuo* to obtain a yellow oil. The crude product was purified by flash chromatography. Elution with dichloromethane afforded 3.40 g of a white solid; 58 % yield; ¹H-NMR (400 MHz; CDCl₃) δ 7.63-7.56 (m, 1H), 7.12-7.04 (m, 1H), 2.812 (s, 3H), 2.807 (s, 3H); ¹⁹F-NMR (376 MHz; CDCl₃) δ -124.91 to -125.03 (m), -127.98 to -128.03 (m), -156.41 to -156.53.

20

Step b: Preparation of 5-dimethylsulfamoyl-2,3,4-trifluoro-benzoic acid

To a cold (-78 °C) stirring solution comprised of 1-dimethylsulfamoyl-2,3,4-trifluorobenzene in anhydrous tetrahydrofuran (60 ml) under a nitrogen atmosphere was added a commercially available lithium diisopropylamide solution (Aldrich, 2.0 M in tetrahydrofuran/heptane/ethylbenzene, 7.5 ml, 0.0150 mol). After stirring for about ten minutes, the purple solution was transferred via canula to a cold, stirring, saturated carbon dioxide in diethyl ether solution (200 ml). The reaction mixture took on a dull burgundy color. The cold bath was removed and the reaction mixture warmed to ambient temperature over one hour. The mixture was then carefully quenched with 10 % aqueous hydrochloric acid (200 ml). The layers were separated. The organic phase was extracted twice (200, 100 ml portions) with 10 % (wt.) aqueous sodium hydroxide. The combined aqueous alkaline extracts were treated with concentrated aqueous hydrochloric acid (100 ml) to pH 0. A white precipitate formed. The suspension was allowed to cool, then was extracted with diethyl ether (600 ml). The organic extract was

35

dried over anhydrous magnesium sulfate and was concentrated *in vacuo* to afford 2.70 g of an off-white solid; 67.5 % yield; mp 225-228 °C; ¹H-NMR (400 MHz; DMSO) δ 14.08 (broad s, 1H), 8.02-7.97 (m, 1H), 2.75 (s, 3H), 2.74 (s, 3H) ¹⁹F-NMR (376 MHz; DMSO) δ -122.50 to -122.63 (m), -122.95 to -123.08 (m), -154.49 to -154.61 (m); MS (APCI+) 284 (M+1, 22), 238 (100); (APCI-) 282 (M-1, 85), 259 (94), 238 (46), 216 (91), 195 (100); IR (KBr) 1702 cm⁻¹; Anal. calcd/found for: C₉H₈F₃NO₄S C, 38.17/38.40; H, 2.85/2.90; N, 4.95/4.80; F, 20.12/19.75; S, 11.32/11.12.

10. Step c: Preparation of 5-dimethylsulfamoyl-2,3,4-trifluoro-benzoic acid methyl ester

The solid 5-dimethylsulfamoyl-2,3,4-trifluoro-benzoic acid (1.47 g, 0.00519 mol) and *p*-toluenesulfonic acid catalyst (17.1 mg) were dissolved in methanol (125 ml). The stirring mixture was brought to reflux under a nitrogen atmosphere for 51 hours. The reaction mixture was concentrated *in vacuo* to give a solid. The product was partitioned between diethyl ether (200 ml) and saturated aqueous potassium carbonate (75 ml). The layers were separated and the organic phase was washed with water (75 ml), brine (75 ml), was dried over anhydrous potassium carbonate, and was concentrated *in vacuo* to afford 0.15 g of an off-white solid; 10 % yield; ¹H-NMR (400 MHz; CDCl₃) δ 8.23-8.19 (m, 1H), 3.92 (s, 3H), 2.83 (s, 6H); ¹⁹F-NMR (376 MHz; CDCl₃) δ -120.79 to -121.02 (m), -153.69 to -153.80.

25. Step d: Preparation of 2-(2-chloro-4-iodo-phenylamino)-5-dimethylsulfamoyl-3,4-difluoro-benzoic acid methyl ester

To a stirring cold (-78 °C) solution comprised of 2-chloro-4-iodoaniline (0.143 g, 5.64x10⁻⁴ mol) in anhydrous tetrahydrofuran (5 ml) under a nitrogen atmosphere was added a commercially available lithium diisopropylamide solution (Aldrich, 2.0 M in tetrahydrofuran/heptane/ethylbenzene, 0.300 ml, 6.0x10⁻⁴ mol). After stirring for 5 minutes, a solution comprised of 5-dimethylsulfamoyl-2,3,4-trifluoro-benzoic acid methyl ester (0.15 g, 5.0x10⁻⁴ mol) in tetrahydrofuran (10 ml) was added via syringe. The cold bath was removed and the reaction mixture was stirred for 2 hours. The reaction mixture was then

partitioned between diethyl ether (125 ml) and saturated aqueous sodium bicarbonate (125 ml). The aqueous bicarbonate phase was extracted with an additional portion (125 ml) of diethyl ether. The combined organic phases were dried over anhydrous magnesium sulfate and concentrated *in vacuo* to give a yellow oil. The oil was crystallized from heptane-ethyl acetate to afford 0.060 g of an off-white powder; 23 % yield; mp 154-156 °C; ¹H-NMR (400 MHz; CDCl₃) δ 9.74 (s, 1H), 8.30 (d, 1H, J=7.1 Hz), 7.72 (s, 1H), 7.49 (d, 1H, J=8.3 Hz), 6.73-6.69 (m, 1H), 3.92 (s, 3H), 2.84 (s, 3H), 2.83 (s, 3H); ¹⁹F-NMR (376 MHz; CDCl₃) δ -123.90 (d), -139.55 (d).

10

EXAMPLE 2

Preparation of 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-sulfamoyl-benzamide (PD 219622)

15

Step a: Preparation of 1-bis-(4-methoxybenzyl)sulfamoyl-2,3,4-trifluorobenzene

To a stirring solution comprised of *bis*-4-methoxybenzylamine (2.5 g, 9.7×10^{-3} mol) and diisopropylethylamine (1.7 ml, 9.7×10^{-3} mol) in dichloromethane (50 ml) at 0 °C under nitrogen atmosphere was added liquid 2,3,4-

trifluorobenzenesulfonyl chloride (2.26 g, 9.5×10^{-3} mol) directly. The mixture was stirred cold for ten minutes. The ice-water bath was removed and the mixture was stirred for an additional 15 minutes and was then diluted with dichloromethane to 350 ml volume and was washed with saturated aqueous ammonium chloride (200 ml). The organic phase was dried (MgSO₄) and concentrated *in vacuo* to afford 4.99 g of a sticky white solid. The crude product was recrystallized from hexanes-acetone to afford 3.00 g of white needles; 70 % yield; mp 87-90 °C; ¹H-NMR (400 MHz; CDCl₃) δ 7.64-7.58 (m, 1H), 7.04-6.99 (m, 1H), [6.97 (d, 4H, J=8.5 Hz), 6.75 (d, 4H, J=8.8 Hz) AB q], 4.33 (s, 4H), 3.76 (s, 6H); ¹⁹F-NMR (376 MHz; CDCl₃) δ -125.44 to -125.56 (m), -128.61 to -128.72 (m), -156.91 to -157.03 (m); MS (APCI+) 121 (M-330, 100); (APCI-) 330 (M-121, 18), 195 (M-256, 100); IR (KBr) 1612, 1517, 1506, 1465, 1258, 1240, 1156, 1037, 1030 cm⁻¹; Anal. calcd/found for: C₂₂H₂₀F₃NO₄S C, 58.53/57.98; H, 4.47/4.61; N, 3.10/2.85.

Step a: Preparation of 5-bis-(4-methoxybenzyl)sulfamoyl-2,3,4-trifluorobenzoic acid

To a stirring solution comprised of 1-bis-(4-methoxybenzyl)sulfamoyl-2,3,4-trifluorobenzene (2.95 g, 6.5×10^{-3} mol) in tetrahydrofuran (60 ml) at -78°C was added a solution comprised of 2.0 M lithium diisopropylamide in tetrahydrofuran/heptane/ethylbenzene (Aldrich, 3.35 ml, 6.7×10^{-3} mol). After several minutes of stirring, the dark solution was transferred via canula over five minutes to a stirring solution comprised of carbon dioxide (excess) in diethyl ether at -78°C . A white precipitate immediately formed. The cold bath was removed and the reaction mixture was stirred at ambient temperature for 18 hours. The reaction mixture was quenched with 200 ml of dilute aqueous hydrochloric acid. The layers were separated and the organic phase was dried (MgSO_4) and concentrated *in vacuo* to give 2.82 g of an off-white solid. Recrystallization from dichloromethane (150 ml) afforded 2.10g of the white powder product; 65 % yield; mp $158-161^\circ\text{C}$; $^1\text{H-NMR}$ (400 MHz; DMSO) δ 7.80-7.76 (m, 1H), 7.05-6.74 (AB q, 8H, $J=8.6$ Hz), 4.33 (s, 4H), 3.66 (s, 6H); $^{19}\text{F-NMR}$ (376 MHz; DMSO) δ -123.28 to -123.36 (m), -124.12 to -124.21 (m), -155.41 to -155.53 (m); MS (APCI-) 494 (M-1, 47), 216 (89), 195 (100); IR (KBr) 3420, 2954, 2838, 1695, 1613, 1512, 1347, 1238, 1152, 1079 cm^{-1} ; Anal. calcd/found for: $\text{C}_{23}\text{H}_{20}\text{F}_3\text{NO}_6\text{S}$ C, 55.76/55.85; H, 4.07/4.02; N, 2.83/2.71; F, 11.50/11.41; S, 6.47/6.25.

Step c: Preparation of 5-bis-(4-methoxybenzyl)sulfamoyl-2-(2-chloro-4-iodophenylamino)-3,4-difluorobenzoic acid (PD 215729)

To a stirring solution comprised of 2-chloro-4-iodoaniline (0.53 g, 2.0×10^{-3} mol) in tetrahydrofuran (10 ml) at -78°C under a nitrogen atmosphere was added a solution comprised of 1.0 M lithium bis(trimethylsilyl)amide in tetrahydrofuran (Aldrich, 4.1 ml, 4.1×10^{-3} mol). Within several minutes the solution became a thick light-green suspension. To this mixture was added a solution comprised of lithium 5-bis-(4-methoxybenzyl)sulfamoyl-2,3,4-trifluorobenzoate in tetrahydrofuran, which was prepared by adding 2.0 ml of the Aldrich lithium bis(trimethylsilyl)amide solution (0.0020 mmol) to a solution comprised of 5-bis-(4-methoxybenzyl)sulfamoyl-2,3,4-trifluorobenzoic acid (1.00 g, 2.0×10^{-3} mol) in

tetrahydrofuran (10 ml) at -78°C . The reaction mixture was stirred for 15 minutes and was then concentrated *in vacuo* to a crude semisolid. The semisolid was taken up into diethyl ether (250 ml) and was washed with 1 % aqueous hydrochloric acid (150 ml). The ether phase was then washed with neutral water (200ml, pH 4 after wash), a second portion of water (200 ml, pH 6 after wash), and brine (200 ml). The organic phase was then dried (MgSO_4) and was concentrated *in vacuo* to give 1.88 g of a sticky residue which was crystallized from toluene-heptane to afford 1.12 g of an off-white powder; 76 % yield; mp $162-166^{\circ}\text{C}$; $^1\text{H-NMR}$ (400 MHz; DMSO) δ 9.86 (s, 1H), 7.92 (d, 1H, $J=6.8$ Hz), 7.86 (d, 1H, $J=1.7$ Hz), 7.60 (dd, 1H, $J=8.5, 1.7$ Hz), 7.06-7.04/6.78-6.75 (AB q, 8H, $J=8.5$ Hz), 6.93-6.89 (m, 1H), 4.31 (s, 4H), 3.66 (s, 6H); $^{19}\text{F-NMR}$ (376 MHz; DMSO) δ -127.22 (d), -141.36 (d); MS (APCI+) 729 ($M+1$, 1), 256 (50), 121 (100); (APCI-) 727 ($M-1$, 100); IR (KBr) 1698, 1673, 1513, 1251 cm^{-1} ; Anal. calcd/found for: $\text{C}_{29}\text{H}_{24}\text{ClF}_2\text{IN}_2\text{O}_6\text{S}$ C, 47.78/47.93; H, 3.32/3.33; N, 3.84/3.80; Cl, 4.86/4.84; F, 5.21/5.46; I, 17.41/17.16; S, 4.40/4.29.

Step d: Preparation of 5-bis-(4-methoxybenzyl)sulfamoyl-2-(2-chloro-4-iodophenylamino)-N-cyclopropylmethoxy-3,4-difluorobenzamide (PD 218774)

To a stirring solution comprised of 5-bis-(4-methoxybenzyl)sulfamoyl-2-(2-chloro-4-iodophenylamino)-3,4-difluorobenzoic acid (0.935 g , $1.28 \times 10^{-3}\text{ mol}$), cyclopropylmethoxylamine hydrochloride (0.175 g , $1.42 \times 10^{-3}\text{ mol}$), and diisopropylethylamine (0.75 ml , $4.26 \times 10^{-3}\text{ mol}$) in a 1:1 v/v tetrahydrofuran-dichloromethane mixture (50 ml) was added solid PyBOP ([benzotriazoloxyl]tripyrrolidino phosphonium hexafluorophosphate, Advanced ChemTech, 0.76 g , $1.46 \times 10^{-3}\text{ mol}$). The reaction mixture was stirred for one hour, was then evaporated to a crude residue which was purified by flash silica column chromatography. Elution with a gradient (25 % dichloromethane to 75 % dichloromethane in hexanes) afforded 0.63 g of the off-white powder product; 62 % yield; mp $70 \rightarrow 300^{\circ}\text{C}$; $^1\text{H-NMR}$ (400 MHz; DMSO) δ 11.92 (s, 1H), 9.35 (s, 1H), 7.60 (s, 1H), 7.50-7.45 (m, 1H), 7.34 (d, 1H, $J=8.5$ Hz), 6.82-6.54 (AB q, 8H, $J=8.3$ Hz), 6.59-6.54 (m, 1H), 4.09 (s, 4H), 3.46 (s, 6H), 0.90-0.80 (m, 1H), 0.30-0.25 (m, 2H), 0.03-0.00 (m, 2H); $^{19}\text{F-NMR}$ (376 MHz; DMSO) δ -129.05 (s), $-$

140.23 (d, J=18.3 Hz); MS (APCI+) 798 (M+1, 70); (APCI-) 796 (M-1, 15), 726 (50), 131 (100); IR (KBr) 1642, 1611, 1584, 1513, 1478 cm^{-1} ; Anal. calcd/found for: $\text{C}_{33}\text{H}_{31}\text{ClF}_2\text{IN}_3\text{O}_6\text{S}$ C, 49.67/49.88; H, 3.92/3.95; N, 5.27/5.19.

5 Step e: Preparation of 2-(2-chloro-4-iodophenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-sulfamoyl-benzamide (PD 219622)

A reaction solution comprised of 5-bis-(4-methoxybenzyl)sulfamoyl-2-(2-chloro-4-iodophenylamino)-N-cyclopropylmethoxy-3,4-difluorobenzamide (0.1010 g, 1.266×10^{-4} mol) in trifluoroacetic acid (4 ml) was stirred at ambient temperature
10 for 24 hours. The mixture was vacuum filtered and the precipitate rinsed with hexanes to afford 28.6 mg of a pale lavender powder; 42 % yield; mp 219-227 °C DEC; $^1\text{H-NMR}$ (400 MHz; DMSO) δ 11.89 (s, 1H), 9.08 (s, 1H), 7.60 (s, 3H), 7.55 (d, 1H, J=6.9 Hz), 7.32 (d, 1H, J=8.6 Hz), 6.63-6.59 (m, 1H), 3.40 (d, 2H, J=6.6 Hz), 0.90-0.80 (m, 1H), 0.30-0.26 (m, 2H), 0.05-0.00 (m, 2H); $^{19}\text{F-NMR}$ (376 MHz;
15 DMSO) δ -130.61 (s), -140.38 (d, J=21.4 Hz); MS (APCI+) 558 (M+1, 70), 282 (100); (APCI-) 556 (M-1, 73), 486 (100); IR (KBr) 3390, 3283, 1652, 1513, 1477, 1163 cm^{-1} ; Anal. calcd/found for: $\text{C}_{17}\text{H}_{15}\text{ClF}_2\text{IN}_3\text{O}_4\text{S} \cdot 0.1 \text{ C}_2\text{HF}_3\text{O}_2$ C, 36.30/36.31; H, 2.67/2.55; N, 7.38/7.00.

20

EXAMPLE 3

Preparation of 2-(2-chloro-4-iodo-phenylamino)-3,4-difluoro-N-hydroxy-5-sulfamoyl-benzamide (PD 224213)

To a stirring solution comprised of 5-bis-(4-methoxybenzyl)sulfamoyl-2-(2-chloro-4-iodophenylamino)-3,4-difluorobenzoic acid (0.67 g, 9.2×10^{-4} mol), O-(tetrahydro-2H-pyran-2-yl)-hydroxylamine (0.113 g, 9.65×10^{-4} mol), and diisopropylethylamine (0.50 ml, 2.9×10^{-3} mol) in a 1:1 v/v tetrahydrofuran-dichloromethane mixture (20 ml) was added solid PyBOP
25 ([benzotriazolyloxy]tripyrrolidino phosphonium hexafluorophosphate, Advanced ChemTech, 0.52 g, 1.0×10^{-3} mol). The reaction mixture was stirred for 30 minutes, was concentrated *in vacuo* to a yellow oil, and was crystallized from methanol to afford 0.35 g of the off-white amorphous intermediate; 46 % yield; the intermediate was dissolved in trifluoroacetic acid (10 ml) and was stirred at ambient temperature for 16 hours. The mixture was vacuum filtered to collect the

precipitate, which was recrystallized from methanol-chloroform to afford 0.055 g of the tan powder product; 26 % yield from intermediate; mp 230-236 °C DEC; ¹H-NMR (400 MHz; DMSO) δ 11.73 (s, 1H), 9.46 (s, 1H), 9.38 (s, 1H), 7.80-7.75 (m, 2H), 7.79 (s, 2H), 7.50 (d, 1H, J=8.5 Hz), 6.82-6.78 (m, 1H); ¹⁹F-NMR (376 MHz; DMSO) δ -130.83 (s), -139.24 (s); MS (APCI+) 504 (M+1, 53), 488 (90), 471 (100); (APCI-) 502 (M-1, 12), 486 (100); IR (KBr) 3295, 1652, 1636, 1519, 1477, 1315, 1157 cm⁻¹; Anal. calcd/found for: C₁₃H₉ClF₂IN₃O₄S · 0.41 CHCl₃ C, 29.15/29.05; H, 1.72/1.66; N, 7.60/7.21.

10

EXAMPLE 4

Preparation of 2-(2-chloro-4-iodo-phenylamino)-3,4-difluoro-5-sulfamoyl-benzoic acid (PD 215730)

Solid 5-*bis*-(4-methoxybenzyl)sulfamoyl-2-(2-chloro-4-iodophenylamino)-3,4-difluorobenzoic acid (0.0995 g, 1.36x10⁻⁴ mol) was dissolved in trifluoroacetic acid (5 ml) under a nitrogen atmosphere. The reaction mixture was stirred at ambient temperature for 65 hours. The mixture was vacuum filtered to isolate 55.2 mg of a fine white precipitate. The crude product was recrystallized from chloroform to afford 31.8 mg of the fluffy white solid product; 48 % yield; mp 295-296 °C DEC; ¹H-NMR (400 MHz; DMSO) δ 9.77 (s, 1H), 8.16 (d, 1H, J=7.3 Hz), 7.82 (s, 3H), 7.56 (d, 1H, J=8.5 Hz), 6.97-6.92 (m, 1H); ¹⁹F-NMR (376 MHz; DMSO) δ -128.47 (s), -141.13 (d, 19.8 Hz); MS (APCI+) 489 (M+1, 5), 102 (100); (APCI-) 491 (32), 490 (18), 489 (100), 488 (18), 487 (M-1, 75); IR (KBr) 3372, 3244, 1688 cm⁻¹; Anal. calcd/found for: C₁₃H₈ClF₂IN₂O₄S C, 31.96/32.19; H, 1.65/1.81; N, 5.73/5.37.

25

EXAMPLE 5

30 Preparation of 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-dimethylsulfamoyl-benzamide (PD 250253)

Step a: Preparation of 2-(2-chloro-4-iodo-phenylamino)-3,4-difluoro-5-dimethylsulfamoyl-benzoic acid (PD 224339)

To a stirring solution comprised of 5-dimethylsulfamoyl-2,3,4-trifluorobenzoic acid (1.00 g, 3.53x10⁻³ mol) in tetrahydrofuran (15 ml) at -78 °C

35

under a nitrogen atmosphere was added a 1.0 M solution of lithium *bis*(trimethylsilyl)amide in tetrahydrofuran (Aldrich, 3.6 ml, 3.6×10^{-3} mol). A lithium 2-chloro-4-iodoanilide suspension formed by adding a 1.0 M solution of lithium *bis*(trimethylsilyl)amide solution (7.2 ml, 7.2×10^{-3} mol) to a solution
5 comprised of 2-chloro-4-iodoaniline (0.94 g, 3.63×10^{-3} mol) in tetrahydrofuran (15 ml) at -78°C was added via canula to the lithium 5-dimethylsulfamoyl-2,3,4-trifluorobenzoate suspension. The cold bath was removed and the reaction mixture was stirred for one hour. The mixture was concentrated *in vacuo* to a crude solid. The crude product was suspended in diethyl ether (200 ml), to which
10 suspension hydrogen chloride gas was introduced to produce a white precipitate. The precipitate was removed by vacuum filtration. The filtrate was concentrated *in vacuo* to give a dull-colored solid, which was triturated with hexanes-dichloromethane to afford 1.31 g of the white powder product; 72 % yield; mp $218-222^\circ\text{C}$; $^1\text{H-NMR}$ (400 MHz; DMSO) δ 9.89 (s, 1H), 8.06 (d, 1H, $J=6.1$ Hz),
15 7.85 (d, 1H, $J=1.9$ Hz), 7.58 (dd, 1H, $J=8.5, 1.9$ Hz), 7.03 (dd, 1H, $J=8.3, 6.6$ Hz), 2.71 (s, 6H); $^{19}\text{F-NMR}$ (376 MHz; DMSO) δ -125.58 (d, $J=18.3$ Hz), -140.14 (d, $J=16.8$ Hz); MS (APCI+) 519 (40), 518 (15), 517 (M+1, 100); (APCI-) 517 (6), 516 (2), 515 (M-1, 5), 480 (45), 127 (100); IR (KBr) 3346, 1665, 1487, 1283 cm^{-1} ; Anal. calcd/found for: $\text{C}_{15}\text{H}_{12}\text{ClF}_2\text{IN}_2\text{O}_4\text{S}$ C, 34.87/34.98; H, 2.34/2.32; N,
20 5.42/5.32.

Step b: Preparation of 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-dimethylsulfamoyl-benzamide

To a suspension comprised of 2-(2-chloro-4-iodo-phenylamino)-3,4-difluoro-5-dimethylsulfamoyl-benzoic acid (0.5 g, 9.68×10^{-4} mol) and
25 cyclopropylmethoxylamine hydrochloride (0.13 g, 1.05×10^{-3} mol) in a 1:1 v/v mixture of dichloromethane-tetrahydrofuran (10 ml) was added diisopropylethylamine (0.65 ml, 3.73×10^{-3} mol) followed by the addition of solid PyBOP (0.55 g, 1.06×10^{-3} mol). The reaction mixture was stirred at ambient temperature for three days. The mixture was concentrated *in vacuo* to a red oil.
30 The crude product was treated with 10 % aqueous hydrochloric acid (150 ml) and was extracted with diethyl ether (150 ml). The organic phase was dried (MgSO_4) and concentrated *in vacuo* to a crude solid. The solid was triturated with

dichloromethane-hexanes and recovered by vacuum filtration to afford 0.3558 g of the white powder product; 63 % yield; mp 222-225 °C DEC; ¹H-NMR (400 MHz; DMSO) δ 11.97 (s, 1H), 9.32 (s, 1H), 7.60 (d, 1H, J=1.9 Hz), 7.49 (d, 1H, J=5.8 Hz), 7.33 (dd, 1H, J=8.4, 1.9 Hz), 6.70 (dd, 1H, 8.4, 6.3 Hz), 3.43 (d, 2H, J=7.2 Hz), 2.53 (s, 6H), 0.87-0.83 (m, 1H), 0.30-0.25 (m, 2H), 0.03-0.00 (m, 2H); ¹⁹F-NMR (376 MHz; DMSO) δ -127.67 (d, J=19.8 Hz), -139.32 (d, J=19.8 Hz); MS (APCI+) 586 (M+1, 100); (APCI-) 584 (M-1, 40), 514 (100); IR (KBr) 3263, 1644, 1585, 1507, 1480 cm⁻¹; Anal. calcd/found for: C₁₉H₁₉ClF₂IN₃O₄S C, 38.96/39.08; H, 3.27/3.18; N, 7.17/7.17.

10

EXAMPLE 6

Preparation of N-cyclopropylmethoxy-3,4-difluoro-5-dimethylsulfamoyl-2-(4-iodo-2-methyl-phenylamino)-benzamide (PD 252745)

15

Step a: Preparation of 3,4-difluoro-5-dimethylsulfamoyl-2-(4-iodo-2-methyl-phenylamino)-benzoic acid (PD 224340)

Same procedure and same scale as Example 4, Step a, except 4-iodo-2-methylaniline was used instead of 2-chloro-4-iodoaniline; afforded 0.9592 g of the off-white powder product; 55 % yield; mp 235-238 °C; ¹H-NMR (400 MHz; DMSO) δ 9.69 (s, 1H), 8.04 (d, 1H, J=6.1 Hz), 7.60 (d, 1H, J=1.5 Hz), 7.45 (dd, 1H, J=8.3, 1.7 Hz), 6.88 (dd, 1H, J=8.3, 5.4 Hz), 2.70 (s, 6H), 2.21 (s, 3H); ¹⁹F-NMR (376 MHz; DMSO) δ -126.25 (d, J=16.8 Hz), -142.74 (d, J=19.8 Hz); MS (APCI+) 497 (M+1, 69), 357 (70), 316 (100); (APCI-) 495 (M-1, 3), 127 (100); IR (KBr) 3240, 1686, 1512, 1473, 1341, 1151 cm⁻¹; Anal. calcd/found for: C₁₆H₁₅F₂IN₂O₄S C, 38.72/38.70; H, 3.05/3.01; N, 5.64/5.49.

25

Step b: Preparation of N-cyclopropylmethoxy-3,4-difluoro-5-dimethylsulfamoyl-2-(4-iodo-2-methyl-phenylamino)-benzamide

30

Same procedure and same scale as Example 4, Step b, except the product was purified by recrystallization from absolute ethanol to afford 0.1718 g of the pale yellow microcrystalline product; 28 % yield; mp 171-172 °C; ¹H-NMR (400 MHz; DMSO) δ 11.79 (s, 1H), 8.91 (s, 1H), 7.40 (d, 1H, J=4.3 Hz), 7.36 (s, 1H), 7.21 (d, 1H, J=8.2 Hz), 6.54 (dd, 1H, 8.2, 4.3 Hz), 3.30 (d, 2H, J=6.5 Hz), 2.52 (s, 6H), 2.00 (s, 3H), 0.85-0.75 (m, 1H), 0.29 (d, 2H, J=7.7 Hz), 0.01 (d, 2H,

35

$J=4.1$ Hz); ^{19}F -NMR (376 MHz; DMSO) δ -128.94 (s), -143.32 (d, $J=19.8$ Hz); MS (APCI+) 566 (M+1, 100); (APCI-) 564 (M-1, 85), 494 (100); IR (KBr) 1649, 1609, 1588, 1512, 1475 cm^{-1} ; Anal. calcd/found for: $\text{C}_{20}\text{H}_{22}\text{F}_2\text{IN}_3\text{O}_4\text{S}$ C, 42.49/42.42; H, 3.92/3.78; N, 7.43/7.40.

5

EXAMPLE 7

Preparation of 2-(2-chloro-4-iodo-phenylamino)-N-hydroxy-4-dimethylsulfamoyl-benzamide

10

Step a: Preparation of 4-methyl-benzene-N,N-dimethylsulfonamide

To a stirring solution comprised of *para*-toluenesulfonyl chloride in dichloromethane at 0 °C is introduced excess gaseous dimethylamine. The precipitate is removed by filtration and the filtrate is concentrated *in vacuo* to obtain the product.

15

Step b: Preparation of 4-methyl-3-nitro-benzene-N,N-dimethylsulfonamide

To a gently stirring solution comprised of 1 molar equivalent of fuming nitric acid in excess concentrated sulfuric acid is added 1 molar equivalent of 4-methyl-benzene-N,N-dimethylsulfonamide in increments. The mixture is stirred for one hour and then poured over chilled water. The mixture is extracted with a suitable solvent like diethyl ether or dichloromethane. The organic phase is dried over a suitable drying agent like magnesium sulfate and concentrated *in vacuo* to afford a crude product which may be purified by normal methods such as chromatography or crystallization from a solvent like chloroform or heptane.

20

25

Step c: Preparation of 3-amino-4-methyl-benzene-N,N-dimethylsulfonamide

The compound 4-methyl-3-nitro-benzene-N,N-dimethylsulfonamide is dissolved in ethanol. A catalyst like Raney nickel is added and the mixture hydrogenated in a shaker. The catalyst is removed by filtration. The solvent is removed *in vacuo* to give a product which may be purified if necessary by chromatography or crystallization from an appropriate solvent like chloroform or heptane-ethyl acetate.

30

Step d: Preparation of 3-fluoro-4-methyl-benzene-N,N-dimethylsulfonamide

The compound 3-amino-4-methyl-benzene-N,N-dimethylsulfonamide is diazotized with an alkyl nitrite like *tert*-butyl nitrite under anhydrous conditions in a non-reactive solvent like tetrahydrofuran or dichloromethane. The intermediate
5 diazonium species is then treated with pyridinium fluoride to give the product, which may be purified by chromatography or crystallization.

Step e: Preparation of 4-dimethylsulfamoyl-2-fluoro-benzoic acid

A mixture comprised of 3-fluoro-4-methyl-benzene-N,N-dimethylsulfonamide and potassium permanganate (2.2 molar equivalents) in
10 water is brought to reflux for four hours. The reaction mixture is filtered through celite. The filtrate is treated with activated carbon and refiltered through fresh celite. The second filtrate is acidified with concentrated hydrochloric acid to pH 0. The mixture is allowed to cool and is extracted with diethyl ether. The organic
15 phase is dried over a drying agent like magnesium sulfate and is concentrated *in vacuo*. The product may be purified by recrystallization from an appropriate solvent like ethanol or chloroform.

Step f: Preparation of 2-(2-chloro-4-iodo-phenylamino)-4-dimethylsulfamoyl-benzoic acid

To a stirring cold (-78 °C) solution comprised of 2-chloro-4-iodoaniline (1 molar equivalent) in anhydrous tetrahydrofuran under a nitrogen atmosphere is added a commercially available lithium diisopropylamide solution (Aldrich, 2.0 M
20 in tetrahydrofuran/heptane/ethylbenzene, 1 molar equivalent). After stirring for 5 minutes, a solution comprised of 4-dimethylsulfamoyl-2-fluoro-benzoic acid (1 molar equivalent) in tetrahydrofuran is added. The cold bath is removed and the reaction mixture is stirred for 2 hours. The reaction mixture is then partitioned between diethyl ether and dilute aqueous hydrochloric acid. The organic phase is washed with brine, dried over magnesium sulfate, and concentrated *in vacuo* to
25 afford a product which may be purified by chromatography of recrystallization from an appropriate solvent like chloroform or heptane-ethanol.
30

Step g: Preparation of 2-(2-chloro-4-iodo-phenylamino)-4-dimethylsulfamoyl-benzoic acid O-(tetrahydro-2H-pyran-2-yl)-oxyamide

A solution comprised of 2-(2-chloro-4-iodo-phenylamino)-4-dimethylsulfamoyl-benzoic acid, O-(tetrahydro-2H-pyran-2-yl)-hydroxylamine (1.25 molar equivalents), benzotriazole-1-yl-oxy-*tris*-pyrrolidino-phosphonium hexafluorophosphate (1.25 molar equivalents), and diisopropylethylamine (3 molar equivalents) in 1:1 v/v tetrahydrofuran-dichloromethane is stirred for 30 minutes. The reaction mixture is concentrated *in vacuo* and the residue is purified by flash chromatography; elution with dichloromethane affords the desired product. The product may be recrystallized with an appropriate solvent like methanol if further purification is necessary.

Step h: Preparation of 2-(2-chloro-4-iodo-phenylamino)-4-dimethylsulfamoyl-N-hydroxy-benzamide

The compound 2-(2-chloro-4-iodo-phenylamino)-4-dimethylsulfamoyl-benzoic acid O-(tetrahydro-2H-pyran-2-yl)-oxyamide is dissolved in an appropriate hydrogen chloride-saturated solvent like methanol or ethanol. Once homogeneous, the solution is concentrated *in vacuo* to give the desired product. The product may be triturated with an appropriate solvent like chloroform or dichloromethane if further purification is necessary.

EXAMPLE 8

Cascade assay for inhibitors of the MAP kinase pathway

Incorporation of ^{32}P into myelin basic protein (MBP) is assayed in the presence of a glutathione S-transferase fusion protein containing p44MAP kinase (GST-MAPK) and a glutathione S-transferase fusion protein containing p45MEK (GST-MEK). The assay solution contains 20 mM HEPES, pH 7.4, 10 mM MgCl_2 , 1 mM MnCl_2 , 1 mM EGTA, 50 μM $[\gamma\text{-}^{32}\text{P}]\text{ATP}$, 10 μg GST-MEK, 0.5 μg GST-MAPK and 40 μg MBP in a final volume of 100 μL . Reactions are stopped after 20 minutes by addition of trichloroacetic acid and filtered through a GF/C filter mat. ^{32}P retained on the filter mat is determined using a 120S Betaplate. Compounds are assessed at 10 μM for ability to inhibit incorporation of ^{32}P .

To ascertain whether compounds are inhibiting GST-MEK or GST MAPK, two additional protocols are employed. In the first protocol, compounds are added to tubes containing GST-MEK, followed by addition of GST-MAPK, MBP and $[\gamma\text{-}^{32}\text{P}]\text{ATP}$. In the second protocol, compounds are added to tubes
5 containing both GST-MEK and GST-MAPK, followed by MBP and $[\gamma\text{-}^{32}\text{P}]\text{ATP}$.

Compounds that show activity in both protocols are scored as MAPK inhibitors, while compounds showing activity in only the first protocol are scored as MEK inhibitors.

10

EXAMPLE 9

In vitro MAP kinase assay

Inhibitory activity can be confirmed in direct assays. For MAP kinase, 1 μg GST-MAPK is incubated with 40 μg MBP for 15 minutes at 30°C in a final volume of 50 μL containing 50 mM Tris (pH 7.5), 10 μM MgCl_2 , 2 μM EGTA, and
15 10 μM $[\gamma\text{-}^{32}\text{P}]\text{ATP}$. The reaction is stopped by addition of Laemmli SDS sample buffer and phosphorylated MBP resolved by electrophoresis on a 10% polyacrylamide gel. Radioactivity incorporated into MBP is determined by both autoradiography, and scintillation counting of excised bands.

20

EXAMPLE 10

In vitro MEK assay

For evaluation of direct MEK activity, 10 μg GST-MEK₁ is incubated with 5 μg of a glutathione S-transferase fusion protein containing p44MAP kinase with a lysine to alanine mutation at position 71 (GST-MAPK-KA). This mutation
25 eliminates kinase activity of MAPK, so only kinase activity attributed to the added MEK remains. Incubations are 15 minutes at 30°C in a final volume of 50 μL containing 50 mM Tris (pH 7.5), 10 μM MgCl_2 , 2 μM EGTA, and 10 μM $[\gamma\text{-}^{32}\text{P}]\text{ATP}$. The reaction is stopped by addition of Laemmli SDS sample buffer. Phosphorylated GST-MAPK-KA is resolved by electrophoresis on a 10%
30 polyacrylamide gel. Radioactivity incorporated into GST-MAPK-KA is determined by autoradiography, and subsequent scintillation counting of excised bands.

Additionally, an artificially activated MEK containing serine to glutamate mutations at positions 218 and 222 (GST-MEK-2E) is used. When these two sites are phosphorylated, MEK activity is increased. Phosphorylation of these sites can be mimicked by mutation of the serine residues to glutamate. For this
5 assay, 5 μ g GST-MEK-2E is incubated with 5 μ g GST-MAPK-KA for 15 minutes at 30°C in the same reaction buffer as described above. Reactions are terminated and analyzed as above.

EXAMPLE 11

10 Whole cell MAP kinase assay

To determine if compounds block activation of MAP kinase in whole cells, the following protocol is used. Cells are plated in multi-well plates and grown to confluence. Cells are serum-deprived overnight. Cells are exposed to the desired concentrations of compound or vehicle (DMSO) for 30 minutes, followed
15 by addition of a growth factor, for example, PDGF (100 ng/mL). After a 5-minute treatment with the growth factor, cells are washed with PBS, and lysed in a buffer consisting of 70 mM NaCl, 10 mM HEPES (pH 7.4), 50 mM glycerol phosphate, and 1% Triton X-100. Lysates are clarified by centrifugation at 13,000 x g for 10 minutes. Five micrograms of the resulting supernatants are incubated with 10 μ g
20 microtubule associated protein-2 (Map2) for 15 minutes at 30°C in a final volume of 25 μ L containing 50 mM Tris (pH 7.4), 10 mM MgCl₂, 2 mM EGTA and 30 μ M [γ -³²P]ATP. Reactions are terminated by addition of Laemmli sample buffer. Phosphorylated Map2 is resolved on 7.5% acrylamide gels and incorporated radioactivity is determined by scintillation counting of excised bands.

25

EXAMPLE 12

Monolayer growth

Cells are plated into multi-well plates at 10 to 20,000 cells/mL. Forty-eight hours after seeding, test compounds are added to the cell growth medium and
30 incubation is continued for 2 additional days. Cells are then removed from the wells by incubation with trypsin and enumerated with a Coulter counter.

EXAMPLE 13

Growth in soft-agar

- 5 Cells are seeded into 35-mm dishes at 5 to 10,000 cells/dish using growth medium containing 0.3% agar. After chilling to solidify the agar, cells are transferred to a 37°C incubator. After 7 to 10 days' growth, visible colonies are manually enumerated with the aid of a dissecting microscope.

EXAMPLE 14

10

Collagen-Induced Arthritis in Mice

- Type II collagen-induced arthritis (CIA) in mice is an experimental model of arthritis that has a number of pathologic, immunologic, and genetic features in common with rheumatoid arthritis. The disease is induced by immunization of
15 DBA/1 mice with 100 µg type II collagen, which is a major component of joint cartilage, delivered intradermally in Freund's complete adjuvant. The disease susceptibility is regulated by the class II MHC gene locus, which is analogous to the association of rheumatoid arthritis with HLA-DR4.

- A progressive and inflammatory arthritis develops in the majority of mice
20 immunized, characterized by paw width increases of up to 100%. A test compound is administered to mice in a range of amounts, such as 20, 60, 100, and 200 mg/kg body weight/day. The duration of the test can be several weeks to a few months, such as 40, 60, or 80 days. A clinical scoring index is used to assess disease progression from erythema and edema (stage 1), joint distortion
25 (stage 2), to joint ankylosis (stage 3). The disease is variable in that it can affect one or all paws in an animal, resulting in a total possible score of 12 for each mouse. Histopathology of an arthritic joint reveals synovitis, pannus formation, and cartilage and bone erosions. All mouse strains that are susceptible to CIA are high antibody responders to type II collagen, and there is a marked cellular
30 response to CII.

EXAMPLE 15

SCW-induced monoarticular arthritis

Arthritis is induced as described by Schwab, *et al.*, *Infection and Immunity*, 59:4436-4442 (1991) with minor modifications. Rats receive 6 µg sonicated SCW
5 [in 10 µl Dulbecco's PBS (DPBS)] by an intraarticular injection into the right tibiotalar joint on day 0. On day 21, the DTH is initiated with 100 µg of SCW (250 µl) administered i.v. For oral compound studies, compounds are suspended in vehicle (0.5% hydroxypropyl-methylcellulose/0.2% Tween 80), sonicated, and administered twice daily (10 ml/kg volume) beginning 1 hr prior to reactivation
10 with SCW. Compounds are administered in amounts between 10 and 500 mg/kg body weight/day, such as 20, 30, 60, 100, 200, and 300 mg/kg/day. Edema measurements are obtained by determining the baseline volumes of the sensitized hindpaw before reactivation on day 21, and comparing them with volumes at subsequent time points such as day 22, 23, 24, and 25. Paw volume
15 is determined by mercury plethysmography.

EXAMPLE 16

Mouse ear-heart transplant model

Fey, T.A. *et al.* describe methods for transplanting split-heart neonatal
20 cardiac grafts into the ear pinna of mice and rats (*J. Pharm. and Toxic. Meth.* 39:9-17 (1998)). Compounds are dissolved in solutions containing combinations of absolute ethanol, 0.2% hydroxypropyl methylcellulose in water, propylene glycol, cremophor, and dextrose, or other solvent or suspending vehicle. Mice are dosed orally or intraperitoneally once, twice or three times daily from the day
25 of transplant (day 0) through day 13 or until grafts have been rejected. Rats are dosed once, twice, or three times daily from day 0 through day 13. Each animal is anesthetized and an incision is made at the base of the recipient ear, cutting only the dorsal epidermis and dermis. The incision is spread open and down to the cartilage parallel to the head, and sufficiently wide to accommodate the
30 appropriate tunneling for a rat or insertion tool for a mouse. A neonatal mouse or rat pup less than 60 hours old is anesthetized and cervically dislocated. The

heart is removed from the chest, rinsed with saline, bisected longitudinally with a scalpel, and rinsed with sterile saline. The donor heart fragment is placed into the preformed tunnel with the insertion tool and air or residual fluid is gently expressed from the tunnel with light pressure. No suturing, adhesive bonding, bandaging, or treatment with antibiotics is required.

Implants are examined at 10-20-fold magnification with a stereoscopic dissecting microscope without anesthesia. Recipients whose grafts are not visibly beating may be anesthetized and evaluated for the presence of electrical activity using Grass E-2 platinum subdermal pin microelectrodes placed either in the pinna or directly into the graft and a tachograph. Implants can be examined 1-4 times a day for 10, 20, 30 or more days. The ability of a test compound to ameliorate symptoms of transplant rejection can be compared with a control compound such as cyclosporine, tacrolimus, or orally-administered leflunomide.

15

EXAMPLE 17

Murine ovalbumin-induced eosinophilia

Female C57BL/6 mice are obtained from the Jackson Laboratory (Bar Harbor, ME). All animals are given food and water ad libitum. Mice are sensitized with a single i.p. injection of OVA (grade V, Sigma Chemical Company, St. Louis, MO) adsorbed to alum, (10 µg OVA + 9 mg alum in 200 µl saline) or vehicle control, (9 mg alum in 200 µl saline) on day 0. On day 14, the mice are challenged with a 12-minute inhalation of an aerosol consisting of 1.5% OVA (weight/volume) in saline produced by a nebulizer (small particle generator, model SPAG-2; ICN Pharmaceuticals, Costa Mesa, CA). Groups of eight mice are dosed with oral vehicle (0.5% hydroxypropylmethylcellulose / 0.25% TWEEN-80), or a test compound at 10, 30, or 100 mg/kg in oral vehicle, 200 µl per mouse p.o. Dosing is performed once per day starting on day 7 or day 13, and extending through day 16.

30

For determination of pulmonary eosinophilia, three days after the first OVA aerosol challenge (day 17), the mice are anesthetized with an i.p. injection of anesthetic (Ketamine/Acepromazine/Xylazine), and the tracheae is exposed

and cannulated. The lungs and upper airways are lavaged twice with 0.5 ml of cold PBS. A portion (200 μ l) of the bronchoalveolar lavage (BAL) fluid is enumerated using a Coulter counter Model ZB1 (Coulter Electronics, Hialeah, FL). The remaining BAL fluid is then centrifuged at 300 x g for five minutes, and
5 the cells are resuspended in 1 ml of HBSS (Gibco BRL) containing 0.5% fetal calf serum (HyClone) and 10 mM HEPES (Gibco BRL). The cell suspension is centrifuged in a cytospin (Shandon Southern Instruments, Sewickley, PA) and stained by Diff Quick (American Scientific Products, McGraw Park, IL) to differentiate BAL leukocytes into neutrophil, eosinophil, monocyte or lymphocyte
10 subsets. The number of eosinophils in the BAL fluid is determined by multiplying the percentage of eosinophils by the total cell count.

EXAMPLE 18

Caco-2 cell studies

15 Cell transport studies were conducted with Caco-2 cells grown on Snapwells between 22 to 28 days postseeding. Typically, 10 mM MES buffer (pH 6.5) with 5 mM KCl, 135 mM NaCl and 1.8 mM CaCl_2 was used for the apical side and 10 mM MOPS (pH 7.4) with 5 mM KCl, 132.5 mM NaCl and 1.8 mM
20 CaCl_2 with 5 mM D-Glucose was used for the basolateral side. After washing the monolayers, appropriate buffers were pipetted into the respective chambers and the cells were pre-equilibrated at 37°C for at least 15 min. On the day of the experiment the growth media was aspirated and the cell monolayers were preequilibrated with appropriate buffers at 37°C for at least 15 min. Thereafter,
25 TEER measurements were performed to confirm the integrity of the monolayers. Transepithelial flux measurements were made by mounting the cell monolayers in a side-by-side diffusion chamber system (Precision Instrument Design, Tahoe City, CA). Temperature was maintained at 37°C with a circulating water jacket. The solutions were mixed with gas-lift circulation with 95% oxygen-5% carbon
30 dioxide. Donor solutions with PD compounds, [^{14}C] mannitol (leakage marker) and [^3H] metoprolol (reference compound) were added to the apical chamber. Donor and receiver samples were collected at selected time intervals for up to 3

hours. Radiolabelled mannitol and metoprolol were analyzed using scintillation counting (TopCount, Packard Instruments, Downers Grove, IL). PD compounds were analyzed using a LC/MS/MS method. Apparent permeability coefficients were calculated using the following equation:

5
$$P_{app} = (V \cdot dC) / (A \cdot C_o \cdot dt)$$

where V= volume of the receiver solution in ml, A = surface area in cm², C_o = initial donor concentration in mM and dC/dt = change in the drug concentration in the receiver chamber over time.

10

EXAMPLE 19

Metabolic Stability in Human and Rat Liver Microsomes

Compounds are individually incubated (5 µM, dissolved in DMSO) with human and rat liver microsomes (0.5 mg/mL protein) in 50 mM KHPO₄ buffer at 37°C in the presence of 1.0 mM NADPH. At 0, 10, 20 and 40 minutes, 100 µL aliquots are removed and added to 300 µL of acetonitrile. Standard curves are run in a similar manner with each compound at concentrations: 7.5 µM, 3.75 µM, 2.5 µM, 1.25 µM. The samples are analyzed for parent concentration by LC/MS/MS. The in vitro metabolic half-life determinations are determined from the concentration vs. time plots using WinNonlin. These in vitro data represent the rate of oxidative and hydrolytic metabolism.

15

20

EXAMPLE 20

Below is a table of selected compounds.

	<u>Compound</u>	<u>Mean IC₅₀</u>	<u>Individual IC₅₀s</u>
5	PD 0298458	0.48	0.56, 0.41
	PD 0298459	0.24	0.16, 0.32
	PD 0298460	0.68	0.54, 0.83
	PD 0298463	5.6	n=1
10	PD 0298464	2.6	2.9, 2.4
	PD 0298465	1.04	1.2, 0.89
	PD 0298467	0.92	1.4, 0.44

15

5979 STRUCTURE CHART

PD Number	APK IC ₅₀ (nM)	C26 Cells (nM)	Structure
193846	222		2-(2-Chloro-4-iodo-phenylamino)-5-dimethylsulfamoyl-3,4-difluoro-benzoic acid methyl ester
215729	3000		5-[Bis-(4-methoxy-benzyl)-sulfamoyl]-2-(2-chloro-4-iodo-phenylamino)-3,4-difluoro-benzoic acid
215730	781		2-(2-Chloro-4-iodo-phenylamino)-3,4-difluoro-5-sulfamoyl-benzoic acid
218774	>3000		5-[Bis-(4-methoxy-benzyl)-sulfamoyl]-2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-benzamide
219622	186	1600	2-(2-Chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-sulfamoyl-benzamide
219795	>3000		2-(2-Chloro-4-iodo-phenylamino)-3,4-difluoro-5-(methyl-pyridin-3-ylmethyl-sulfamoyl)-benzoic acid
224213	122	9100	2-(2-Chloro-4-iodo-phenylamino)-3,4-difluoro-N-hydroxy-5-sulfamoyl-benzamide
224339	283.5	>10,000	2-(2-Chloro-4-iodo-phenylamino)-5-dimethylsulfamoyl-3,4-difluoro-benzoic acid
224340	440.5	>10,000	5-Dimethylsulfamoyl-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzoic acid
250253	936.5		2-(2-Chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-5-dimethylsulfamoyl-3,4-difluoro-benzamide
252745	3700		N-Cyclopropylmethoxy-5-dimethylsulfamoyl-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide


F. OTHER EMBODIMENTS

From the above disclosure and examples, and from the claims below, the
5 essential features of the invention are readily apparent. The scope of the
invention also encompasses various modifications and adaptations within the
knowledge of a person of ordinary skill. Examples include a disclosed compound
modified by addition or removal of a protecting group, or an ester, pharmaceutical
salt, hydrate, acid, or amide of a disclosed compound. Publications cited herein
10 are hereby incorporated by reference in their entirety.

What is claimed is:

CLAIMS

1. A compound of formula (I):



(I)

wherein

W is OR_1 , NR_2OR_1 , NR_AR_B , $NR_2NR_AR_B$, or $NR_2(CH_2)_{2-4}NR_AR_B$;

- 20 R_1 is H, C₁₋₈ alkyl, C₃₋₈ alkenyl, C₃₋₈ alkynyl, C₃₋₈ cycloalkyl, phenyl, (phenyl)C₁₋₄ alkyl, (phenyl)C₃₋₄ alkenyl, (phenyl)C₃₋₄ alkynyl, (C₃₋₈ cycloalkyl)-C₁₋₄ alkyl, (C₃₋₈ cycloalkyl)C₃₋₄ alkenyl, (C₃₋₈ cycloalkyl)C₃₋₄ alkynyl, C₃₋₈ heterocyclic radical, (C₃₋₈ heterocyclic radical)C₁₋₄ alkyl, (C₃₋₈ heterocyclic radical)C₃₋₄ alkenyl, (C₃₋₈ heterocyclic radical)C₃₋₄ alkynyl or $(CH_2)_{2-4}NR_AR_B$;

25 R_2 is H, phenyl, C₁₋₄ alkyl, C₃₋₄ alkenyl, C₃₋₈ alkynyl, C₃₋₈ cycloalkyl, or (C₃₋₈ cycloalkyl)-C₁₋₄ alkyl;

- 30 R_A is H, C₁₋₆ alkyl, C₃₋₈ alkenyl, C₃₋₈ alkynyl, C₃₋₈ cycloalkyl, phenyl, (C₃₋₈ cycloalkyl)C₁₋₄ alkyl, (C₃₋₈ cycloalkyl)C₃₋₄ alkenyl, (C₃₋₈ cycloalkyl)C₃₋₄ alkynyl, C₃₋₈ heterocyclic radical, (C₃₋₈ heterocyclic radical)C₁₋₄ alkyl, (aminosulfonyl)phenyl, [(aminosulfonyl)phenyl]C₁₋₄ alkyl, (aminosulfonyl)C₁₋₆ alkyl, (aminosulfonyl)C₃₋₆ cycloalkyl, or [(aminosulfonyl)C₃₋₆ cycloalkyl]C₁₋₄ alkyl;

R_B is H, C₁₋₈ alkyl, C₃₋₈ alkenyl, C₃₋₈ alkynyl, C₃₋₈ cycloalkyl, or C₆₋₈ aryl;

R_3 is H, F, Cl, Br, or NO₂;

5

R_4 is H or F;

R_5 is H, methyl or Cl;

10 R_6 is H, C₁₋₄ alkyl, hydroxyethyl, hydroxypropyl, (CH₂)₂₋₄(NR_CR_D), phenyl, 2-pyridyl, 3-pyridyl, 4-pyridyl or CH₂Ar, where Ar is phenyl, 2-pyridyl, 3-pyridyl, or 4-pyridyl;

15 R_7 is H, C₁₋₄ alkyl, hydroxyethyl, hydroxypropyl, (CH₂)₂₋₄(NR_CR_D), phenyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, or CH₂Ar', where Ar' is phenyl, 2-pyridyl, 3-pyridyl, or 4-pyridyl;

20 each of R_C and R_D is independently selected from H, C₁₋₆ alkyl, C₃₋₄ alkenyl, C₃₋₄ alkynyl, C₃₋₆ cycloalkyl, C₃₋₆ heterocyclic radical, and phenyl; NR_CR_D can also be selected from morpholinyl, piperazinyl, pyrrolidinyl, or piperadiny;

25 wherein each hydrocarbon radical or heterocyclic radical above is optionally substituted with between 1 and 3 substituents independently selected from halo, C₁₋₄ alkyl, C₃₋₆ cycloalkyl, C₂₋₄ alkenyl, C₂₋₄ alkynyl, phenyl, hydroxy, amino, (amino)sulfonyl, and NO₂, wherein each substituent alkyl, cycloalkyl, alkenyl, alkynyl or phenyl is in turn optionally substituted with between 1 and 3 substituents independently selected from halo, C₁₋₂ alkyl, hydroxy, amino, and NO₂;

30 or a pharmaceutically-acceptable salt or C₁₋₆ ester thereof.

2. A compound of claim 1, wherein the sulfamoyl group is *meta* to W(CO)- and *para* to the bridging NH.
3. A compound of claim 1, wherein the sulfamoyl group is *para* to W(CO)- and *meta* to the bridging NH.
4. A compound of claim 1, wherein R₄ is fluoro.
5. A compound of claim 1, where R₃ is fluoro.
6. A compound of claim 1, where R₃ is H.
7. A compound of claim 1, wherein W is OH.
8. A compound of claim 1, wherein W is NR₂OR₁.
9. A compound of claim 7, wherein each of R₃ and R₄ is fluoro.
10. A compound of claim 1, wherein R₁ has at least one hydroxy substituent.
11. A compound of claim 1, wherein R₁ is H, methyl, ethyl, propyl, isopropyl, isobutyl, benzyl, phenethyl, allyl, C₃₋₅ alkenyl, C₃₋₅ alkynyl, C₃₋₆ cycloalkyl, (C₃₋₅ cycloalkyl)C₁₋₂ alkyl, or (C₃₋₅ heterocyclic radical)-C₁₋₂ alkyl.
12. A compound of claim 11, wherein R₁ is H or (C₃₋₄ cycloalkyl)-C₁₋₂ alkyl.
13. A compound of claim 1, wherein R₂ is H, methyl, C₃₋₄ alkynyl, C₃₋₅ cycloalkyl, or (C₃₋₅ cycloalkyl)methyl.

14. A compound of claim 1, wherein R_A is H, methyl, ethyl, isobutyl, hydroxyethyl, hydroxypropyl, cyclopropylmethyl, cyclobutylmethyl, C₃₋₄ alkynyl, phenyl, 2-piperidin-1-yl-ethyl, 2,3-dihydroxy-propyl, 3-[4-(2-hydroxyethyl)-piperazin-1-yl]-propyl, 2-pyrrolidin-1-yl-ethyl, or 2-diethylamino-ethyl; and R_B is H; or where R_B is methyl and R_A is phenyl.

15. A compound of claim 1, wherein R₇ is (CH₂)₂₋₄(NR_CR_D).

16. A compound of claim 1, wherein $\text{NR}_\text{C}\text{R}_\text{D}$ is selected from morpholinyl, piperazinyl, pyrrolidinyl, or piperadinyl.

17. A compound of claim 1, wherein R₅ is methyl or chloro.

18. A compound of claim 1 having the structure: 2-(2-chloro-4-iodo-phenylamino)-4-sulfamoyl-benzoic acid; 2-(2-chloro-4-iodo-phenylamino)-N-hydroxy-4-sulfamoyl-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-4-sulfamoyl-benzamide; 2-(2-chloro-4-iodo-phenylamino)-4-(2-morpholin-4-yl-ethylsulfamoyl)-benzoic acid; 2-(2-chloro-4-iodo-phenylamino)-N-hydroxy-4-(2-morpholin-4-yl-ethylsulfamoyl)-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-4-(2-morpholin-4-yl-ethylsulfamoyl)-benzamide; 2-(2-chloro-4-iodo-phenylamino)-3,4-difluoro-5-sulfamoyl-benzoic acid; 2-(2-chloro-4-iodo-phenylamino)-3,4-difluoro-N-hydroxy-5-sulfamoyl-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-sulfamoyl-benzamide; 2-(2-chloro-4-iodo-phenylamino)-3,4-difluoro-5-(2-morpholin-4-yl-ethylsulfamoyl)-benzoic acid; 2-(2-chloro-4-iodo-phenylamino)-3,4-difluoro-N-hydroxy-5-(2-morpholin-4-yl-ethylsulfamoyl)-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-(2-morpholin-4-yl-ethylsulfamoyl)-benzamide; 5-(bis-pyridin-3-ylmethyl-sulfamoyl)-3,4-difluoro-2-(4-iodo-phenylamino)-benzoic acid; 5-(bis-pyridin-3-ylmethyl-sulfamoyl)-N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-phenylamino)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-phenylamino)-5-(methyl-pyridin-3-ylmethyl-sulfamoyl)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-

phenylamino)-5-[(pyridin-3-ylmethyl)-sulfamoyl]-benzamide; N-cyclopropylmethoxy-5-[(3-diethylamino-propyl)-pyridin-3-ylmethyl-sulfamoyl]-3,4-difluoro-2-(4-iodo-phenylamino)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-5-[(3-hydroxy-propyl)-pyridin-3-ylmethyl-sulfamoyl]-2-(4-iodo-phenylamino)-benzamide; N-cyclopropylmethoxy-5-(ethyl-pyridin-3-ylmethyl-sulfamoyl)-3,4-difluoro-2-(4-iodo-phenylamino)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-5-[(2-hydroxy-ethyl)-pyridin-3-ylmethyl-sulfamoyl]-2-(4-iodo-phenylamino)-benzamide; 5-(bis-pyridin-2-ylmethyl-sulfamoyl)-3,4-difluoro-2-(4-iodo-phenylamino)-benzoic acid; 5-(bis-pyridin-2-ylmethyl-sulfamoyl)-N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-phenylamino)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-phenylamino)-5-(methyl-pyridin-2-ylmethyl-sulfamoyl)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-phenylamino)-5-[(pyridin-2-ylmethyl)-sulfamoyl]-benzamide; 5-(bis-pyridin-3-ylmethyl-sulfamoyl)-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzoic acid; 5-(bis-pyridin-3-ylmethyl-sulfamoyl)-N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-5-(methyl-pyridin-3-ylmethyl-sulfamoyl)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-5-[(pyridin-3-ylmethyl)-sulfamoyl]-benzamide; N-cyclopropylmethoxy-5-[(3-diethylamino-propyl)-pyridin-3-ylmethyl-sulfamoyl]-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-5-[(3-hydroxy-propyl)-pyridin-3-ylmethyl-sulfamoyl]-2-(4-iodo-2-methyl-phenylamino)-benzamide; N-cyclopropylmethoxy-5-(ethyl-pyridin-3-ylmethyl-sulfamoyl)-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-5-[(2-hydroxy-ethyl)-pyridin-3-ylmethyl-sulfamoyl]-2-(4-iodo-2-methyl-phenylamino)-benzamide; 5-(bis-pyridin-2-ylmethyl-sulfamoyl)-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzoic acid; 5-(bis-pyridin-2-ylmethyl-sulfamoyl)-N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-5-(methyl-pyridin-2-ylmethyl-sulfamoyl)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-5-[(pyridin-2-ylmethyl)-sulfamoyl]-benzamide; 5-(bis-pyridin-3-ylmethyl-sulfamoyl)-2-(2-chloro-4-iodo-phenylamino)-3,4-difluoro-

benzoic acid; 5-(bis-pyridin-3-ylmethyl-sulfamoyl)-2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-(methyl-pyridin-3-ylmethyl-sulfamoyl)-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-[(pyridin-3-ylmethyl)-sulfamoyl]-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-5-[(3-diethylamino-propyl)-pyridin-3-ylmethyl-sulfamoyl]-3,4-difluoro-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-[(3-hydroxy-propyl)-pyridin-3-ylmethyl-sulfamoyl]-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-5-(ethyl-pyridin-3-ylmethyl-sulfamoyl)-3,4-difluoro-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-[(2-hydroxy-ethyl)-pyridin-3-ylmethyl-sulfamoyl]-benzamide; 5-(bis-pyridin-2-ylmethyl-sulfamoyl)-2-(2-chloro-4-iodo-phenylamino)-3,4-difluoro-benzoic acid; 5-(bis-pyridin-2-ylmethyl-sulfamoyl)-2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-(methyl-pyridin-2-ylmethyl-sulfamoyl)-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-[(pyridin-2-ylmethyl)-sulfamoyl]-benzamide; N-cyclopropylmethoxy-3,4-difluoro-5-[(3-hydroxy-propyl)-pyridin-2-ylmethyl-sulfamoyl]-2-(4-iodo-phenylamino)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-5-[(2-hydroxy-ethyl)-pyridin-2-ylmethyl-sulfamoyl]-2-(4-iodo-phenylamino)-benzamide; 5-(benzyl-pyridin-2-ylmethyl-sulfamoyl)-N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-phenylamino)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-phenylamino)-5-[(pyridin-4-ylmethyl)-sulfamoyl]-benzamide; N-cyclopropylmethoxy-5-(ethyl-pyridin-4-ylmethyl-sulfamoyl)-3,4-difluoro-2-(4-iodo-phenylamino)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-phenylamino)-5-(methyl-pyridin-4-ylmethyl-sulfamoyl)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-5-[(3-hydroxy-propyl)-pyridin-4-ylmethyl-sulfamoyl]-2-(4-iodo-phenylamino)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-5-[(2-hydroxy-ethyl)-pyridin-4-ylmethyl-sulfamoyl]-2-(4-iodo-phenylamino)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-phenylamino)-5-(methyl-phenyl-sulfamoyl)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-phenylamino)-5-phenylsulfamoyl-

benzamide; N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-phenylamino)-5-(pyridin-3-ylsulfamoyl)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-5-[(3-hydroxy-propyl)-pyridin-2-ylmethyl-sulfamoyl]-2-(4-iodo-2-methyl-phenylamino)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-5-[(2-hydroxy-ethyl)-pyridin-2-ylmethyl-sulfamoyl]-2-(4-iodo-2-methyl-phenylamino)-benzamide; 5-(benzyl-pyridin-2-ylmethyl-sulfamoyl)-N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-5-[(pyridin-4-ylmethyl)-sulfamoyl]-benzamide; N-cyclopropylmethoxy-5-(ethyl-pyridin-4-ylmethyl-sulfamoyl)-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-5-(methyl-pyridin-4-ylmethyl-sulfamoyl)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-5-[(3-hydroxy-propyl)-pyridin-4-ylmethyl-sulfamoyl]-2-(4-iodo-2-methyl-phenylamino)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-5-[(2-hydroxy-ethyl)-pyridin-4-ylmethyl-sulfamoyl]-2-(4-iodo-2-methyl-phenylamino)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-5-(methyl-phenyl-sulfamoyl)-benzamide; N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-5-phenylsulfamoyl-benzamide; N-cyclopropylmethoxy-3,4-difluoro-2-(4-iodo-2-methyl-phenylamino)-5-(pyridin-3-ylsulfamoyl)-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-[(3-hydroxy-propyl)-pyridin-2-ylmethyl-sulfamoyl]-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-[(2-hydroxy-ethyl)-pyridin-2-ylmethyl-sulfamoyl]-benzamide; 5-(benzyl-pyridin-2-ylmethyl-sulfamoyl)-2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-[(pyridin-4-ylmethyl)-sulfamoyl]-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-5-(ethyl-pyridin-4-ylmethyl-sulfamoyl)-3,4-difluoro-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-(methyl-pyridin-4-ylmethyl-sulfamoyl)-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-[(3-hydroxy-propyl)-pyridin-4-ylmethyl-sulfamoyl]-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-[(2-hydroxy-ethyl)-pyridin-4-ylmethyl-sulfamoyl]-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-

- 3,4-difluoro-5-(methyl-phenyl-sulfamoyl)-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-phenylsulfamoyl-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-3,4-difluoro-5-(pyridin-3-ylsulfamoyl)-benzamide; N-cyclopropylmethoxy-2-(4-iodo-phenylamino)-4-phenylsulfamoyl-benzamide; N-cyclopropylmethoxy-2-(4-iodo-phenylamino)-4-(pyridin-3-ylsulfamoyl)-benzamide; N-cyclopropylmethoxy-2-(4-iodo-phenylamino)-4-[(pyridin-3-ylmethyl)-sulfamoyl]-benzamide; 4-(bis-pyridin-3-ylmethyl-sulfamoyl)-N-cyclopropylmethoxy-2-(4-iodo-phenylamino)-benzamide; N-cyclopropylmethoxy-4-[(2-hydroxy-ethyl)-pyridin-4-ylmethyl-sulfamoyl]-2-(4-iodo-phenylamino)-benzamide; N-cyclopropylmethoxy-2-(4-iodo-phenylamino)-4-(methyl-pyridin-3-ylmethyl-sulfamoyl)-benzamide; N-cyclopropylmethoxy-4-[(3-diethylamino-propyl)-pyridin-3-ylmethyl-sulfamoyl]-2-(4-iodo-phenylamino)-benzamide; N-cyclopropylmethoxy-2-(4-iodo-2-methyl-phenylamino)-4-phenylsulfamoyl-benzamide; N-cyclopropylmethoxy-2-(4-iodo-2-methyl-phenylamino)-4-(pyridin-3-ylsulfamoyl)-benzamide; N-cyclopropylmethoxy-2-(4-iodo-2-methyl-phenylamino)-4-[(pyridin-3-ylmethyl)-sulfamoyl]-benzamide; 4-(bis-pyridin-3-ylmethyl-sulfamoyl)-N-cyclopropylmethoxy-2-(4-iodo-2-methyl-phenylamino)-benzamide; N-cyclopropylmethoxy-4-[(2-hydroxy-ethyl)-pyridin-4-ylmethyl-sulfamoyl]-2-(4-iodo-2-methyl-phenylamino)-benzamide; N-cyclopropylmethoxy-2-(4-iodo-2-methyl-phenylamino)-4-(methyl-pyridin-3-ylmethyl-sulfamoyl)-benzamide; N-cyclopropylmethoxy-4-[(3-diethylamino-propyl)-pyridin-3-ylmethyl-sulfamoyl]-2-(4-iodo-2-methyl-phenylamino)-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-4-phenylsulfamoyl-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-4-(pyridin-3-ylsulfamoyl)-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-4-[(pyridin-3-ylmethyl)-sulfamoyl]-benzamide; 4-(bis-pyridin-3-ylmethyl-sulfamoyl)-2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-4-[(2-hydroxy-ethyl)-pyridin-4-ylmethyl-sulfamoyl]-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-4-(methyl-pyridin-3-ylmethyl-sulfamoyl)-benzamide; 2-(2-chloro-4-iodo-phenylamino)-N-cyclopropylmethoxy-4-[(3-diethylamino-propyl)-pyridin-3-ylmethyl-sulfamoyl]-

benzamide; 5-[bis-(4-methoxy-benzyl)-sulfamoyl]-2-(2-chloro-4-iodo-phenylamino)-3,4-difluoro-benzoic acid; and 2-(2-chloro-4-iodo-phenylamino)-5-dimethylsulfamoyl-3,4-difluoro-benzoic acid methyl ester.

- 5 19. A compound of claim 1, having a structure selected from:
PD 298458, N-Allyloxy-2-(2-chloro-4-iodo-phenylamino)-3,4-difluoro-5-(4-methyl-piperazine-1-sulfonyl)-benzamide; PD 298459, N-Allyloxy-2-(2-chloro-4-iodo-phenylamino)-3,4-difluoro-5-(methyl-phenyl-sulfamoyl)-benzamide; PD
298460, 5-(Allyl-methyl-sulfamoyl)-N-allyloxy-2-(2-chloro-4-iodo-phenylamino)-
10 3,4-difluoro-benzamide; PD 298463, 1-[5-Allyloxycarbamoyl-4-(2-chloro-4-iodo-phenylamino)-2,3-difluoro-benzenesulfonyl]-piperidine-3-carboxylic acid amide;
PD 298464, N-Allyloxy-2-(2-chloro-4-iodo-phenylamino)-5-[(3-dimethylamino-propyl)-methyl-sulfamoyl]-3,4-difluoro-benzamide; PD 298465, N-Allyloxy-2-(2-chloro-4-iodo-phenylamino)-3,4-difluoro-5-(4-pyridin-2-yl-piperazine-1-sulfonyl)-
15 benzamide; and PD 298467, N-Allyloxy-2-(2-chloro-4-iodo-phenylamino)-3,4-difluoro-5-(methoxy-methyl-sulfamoyl)-benzamide.

20. A pharmaceutical composition comprising a compound of claim 1 and a pharmaceutically-acceptable carrier.

20

21. A method for treating a proliferative disease, said method comprising administering to a patient in need of such treatment a pharmaceutically-effective amount of a composition comprising a compound of claim 1.

25

22. A method of claim 20, wherein said proliferative disease is selected from psoriasis, restenosis, autoimmune disease, and atherosclerosis.

23. A method for treating cancer, said method comprising administering
30 to a patient in need of such treatment a pharmaceutically-effective amount of a composition comprising a compound of claim 1.

24. A method of claim 23, wherein said cancer is MEK-related.

25. A method of claim 23, wherein said cancer is brain, breast, lung, ovarian, pancreatic, prostate, renal, or colorectal cancer.

5

26. A method for treating, or ameliorating the sequelae of, a stroke, said method comprising administering to a patient in need of such treatment a pharmaceutically-effective amount of a composition comprising a compound of claim 1.

10

27. A method for treating, or ameliorating the sequelae of, heart failure, said method comprising administering to a patient in need of such treatment a pharmaceutically-effective amount of a composition comprising a compound of claim 1.

15

28. A method for treating or reducing the symptoms of xenograft rejection, said method comprising administering to a cell(s) transplant, limb transplant, skin transplant, organ transplant or bone marrow transplant patient a pharmaceutically-effective amount of a composition comprising a compound of claim 1.

20

29. A method for treating osteoarthritis, said method comprising administering to a patient in need of such treatment a pharmaceutically-effective amount of a composition comprising a compound of claim 1.

25

30. A method for treating rheumatoid arthritis, said method comprising administering to a patient in need of such treatment a pharmaceutically-effective amount of a composition comprising a compound of claim 1.

30

31. A method for treating cystic fibrosis, said method comprising administering to a patient in need of such treatment a pharmaceutically-effective amount of a composition comprising a compound of claim 1.

32. A method for treating hepatomegaly, said method comprising administering to a patient in need of such treatment a pharmaceutically-effective amount of a composition comprising a compound of claim 1.

5

33. A method for treating cardiomegaly, said method comprising administering to a patient in need of such treatment a pharmaceutically-effective amount of a composition comprising a compound of claim 1.

10

34. A method for treating Alzheimer's disease, said method comprising administering to a patient in need of such treatment a pharmaceutically-effective amount of a composition comprising a compound of claim 1.

15

35. A method for treating a complication of diabetes, said method comprising administering to a patient in need of such treatment a pharmaceutically-effective amount of a composition comprising a compound of claim 1.

20

36. A method for treating septic shock, said method comprising administering to a patient in need of such treatment a pharmaceutically-effective amount of a composition comprising a compound of claim 1.

25

37. A method for treating a viral infection, said method comprising administering to a patient in need of such treatment a pharmaceutically-effective amount of a composition comprising a compound of claim 1.

38. A method of claim 36, wherein said viral infection is an infection of HIV.

30

39. A method for treating cancer, said method comprising (a) administering to a patient in need of such treatment, a pharmaceutically-effective amount of a composition comprising a compound of claim 1; and

(b) administering a therapy selected from radiation therapy and chemotherapy.

40. A method of claim 39, wherein said chemotherapy comprises a mitotic inhibitor.

5

41. A method of claim 40, wherein said mitotic inhibitor is selected from paclitaxel, docetaxel, vincristine, vinblastine, vinorelbine, and vinflunine.

10

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/30435

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07C311/39 A61K31/18 A61P5/48 A61P17/06 A61P19/02
A61P25/28 A61P31/18 A61P35/00 A61P37/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

-IPC 7 C07C A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 98 37881 A (WARNER LAMBERT) 3 September 1998 (1998-09-03) page 3 -page 4	1-41
A	US 5 525 625 A (A.J. BRIDGES, ET AL.) 11 June 1996 (1996-06-11) cited in the application the whole document	1-41

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

12 May 2000

Date of mailing of the international search report

26/05/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

English, R

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/ 30435

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 21-41
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 21-41
are directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int: Jonal Application No

PCT/US 99/30435

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9837881 A	03-09-1998	AU 5610398 A ZA 9801578 A	18-09-1998 02-09-1998
US 5525625 A	11-06-1996	AT 181913 T AU 690400 B AU 4245696 A CA 2208075 A DE 69510696 D DE 69510696 T EP 0805807 A ES 2137554 T GR 3031295 T JP 10512878 T NZ 297320 A WO 9622985 A ZA 9600528 A	15-07-1999 23-04-1998 14-08-1996 01-08-1996 12-08-1999 23-12-1999 12-11-1997 16-12-1999 31-12-1999 08-12-1998 28-05-1999 01-08-1996 15-08-1996

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☒ **FADED TEXT OR DRAWING**
- ☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☒ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.